

# The method of the weakly conjugate operator: Extensions and applications to operators on graphs and groups

M. Măntoiu<sup>1</sup>, S. Richard<sup>2\*</sup> and R. Tiedra de Aldecoa<sup>3</sup>

<sup>1</sup> *Departamento de Matemáticas, Universidad de Chile, Las Palmeras 3425, Casilla 653, Santiago, Chile*

<sup>2</sup> *Department of Pure Mathematics and Mathematical Statistics, Centre for Mathematical Sciences, University of Cambridge, Cambridge, CB3 0WB, United Kingdom*

<sup>3</sup> *CNRS (UMR 8088) and Department of Mathematics, University of Cergy-Pontoise, 2 avenue Adolphe Chauvin, 95302 Cergy-Pontoise Cedex, France*

E-mails: *Marius.Mantoiu@imar.ro, sr510@cam.ac.uk, rafael.tiedra@u-cergy.fr*

## Abstract

In this review we present some recent extensions of the method of the weakly conjugate operator. We illustrate these developments through examples of operators on graphs and groups.

## 1 Introduction

In spectral analysis, one of the most powerful tools is the method of the conjugate operator, also called Mourre's commutator method after the seminal work of Mourre in the early eighties. This approach has reached a very high degree of precision and abstraction in [1]; see also [14] for further developments. In order to study the nature of the spectrum of a selfadjoint operator  $H$ , the main idea of the standard method is to find an auxiliary selfadjoint operator  $A$  such that the commutator  $i[H, A]$  is strictly positive when localized in some interval of the spectrum of  $H$ . More precisely, one looks for intervals  $J$  of  $\mathbb{R}$  such that

$$E(J)i[H, A]E(J) \geq aE(J) \quad (1.1)$$

for some strictly positive constant  $a$  that depends on  $J$ , where  $E(J)$  denotes the spectral projection of  $H$  on the interval  $J$ . An additional compact contribution to (1.1) is allowed, greatly enlarging the range of applications.

When strict positivity is not available, one can instead look for an  $A$  such that the commutator is positive and injective, *i.e.*

$$i[H, A] > 0. \quad (1.2)$$

This requirement is close to the one of the Kato-Putnam theorem, *cf.* [29, Thm. XIII.28]. A new commutator method based on such an inequality was proposed in [7, 8]. By analogy to the method of the conjugate operator, it has been called the method of the weakly conjugate operator (MWCO). Under some technical assumptions, both approaches lead to a limiting absorption principle, that is, a control of the resolvent of  $H$  near the real axis. In the case of the usual method of the conjugate operator, this result is obtained locally in  $J$ , and away from thresholds. The MWCO establishes the existence of the boundary value of resolvent also at thresholds, but originally applies only to situations where the operator  $H$  has a purely absolutely continuous spectrum. This drawback limits drastically the range of applications. However, in some recent works [26, 27, 30] the MWCO has also been applied successfully to examples with point spectrum. This review intend to present and illustrate some of these extensions through applications to the spectral theory of operators acting on groups and graphs.

---

\*On leave from Université de Lyon; Université Lyon 1; INSA de Lyon, F-69621; Ecole Centrale de Lyon; CNRS, UMR5208, Institut Camille Jordan, 43 blvd du 11 novembre 1918, F-69622 Villeurbanne-Cedex, France.

Compared to the huge number of applications based on an inequality of the form (1.1), the number of papers that contain applications of the MWCO is very small. Let us cite for example the works [13, 24, 25] that deal with the original form of the theory, and the papers [12, 20] that contain very close arguments. The derivation of the limiting absorption principle of [12] has been abstracted in [30]. The framework of [30] is still the one of the MWCO, but since its result applies a certain class of two-body Schrödinger operators which have bound states below zero, it can also be considered as the first extension of the MWCO dealing with operators that are not purely absolutely continuous. The main idea of [30] is that  $H$  itself can add some positivity to (1.2). The new requirement is the existence of a constant  $c \geq 0$  such that

$$-cH + i[H, A] > 0.$$

This inequality, together with some technical assumptions, lead to a limiting absorption principle which is either uniform on  $\mathbb{R}$  if  $c = 0$  or uniform on  $[0, \infty)$  if  $c > 0$ .

The extensions of the MWCO developed in [26, 27] are of a different nature. In these papers, the operators  $H$  under consideration admit a natural conjugate operator  $A$  that fulfills the inequality  $i[H, A] \geq 0$ , namely, the commutator is positive but injectivity may fail. In that situation, the authors considered a decomposition  $\mathcal{H} := \mathcal{K} \oplus \mathcal{G}$ , and the restrictions of  $H$  and  $i[H, A]$  to these subspaces. In favorable circumstances the injectivity can be restored in one of the subspace, and a comprehensible description of the vectors of the second subspace can be given. This decomposition leads again to statements that are close to the ones of the MWCO, but which apply to operators with arbitrary spectrum. This extension is described in Section 2.2.

We would also like to mention the references [2, 3]. They perform what can be considered as a unitary version of the MWCO and extend the Kato-Putnam analysis of unitary operators to the case of unbounded conjugate operators. The main applications concern time-depending propagators.

The content of this review paper is the following. In Section 2 we recall the original method of the weakly conjugate operator, and then present an abstract version of the approach used in [26, 27]. In Section 3 we present applications of this approach to the study of adjacency operators on graphs. A similar analysis for operators of convolution on locally compact groups is performed in Section 4.

Let us finally fix some notations. Given a selfadjoint operator  $H$  in a Hilbert space  $\mathcal{H}$ , we write  $\mathcal{H}_c(H)$ ,  $\mathcal{H}_{ac}(H)$ ,  $\mathcal{H}_{sc}(H)$ ,  $\mathcal{H}_s(H)$  and  $\mathcal{H}_p(H)$  respectively for the continuous, absolutely continuous, singularly continuous, singular and pure point subspaces of  $\mathcal{H}$  with respect to  $H$ . The corresponding parts of the spectrum of  $H$  are denoted by  $\sigma_c(H)$ ,  $\sigma_{ac}(H)$ ,  $\sigma_{sc}(H)$ ,  $\sigma_s(H)$  and  $\sigma_p(H)$ .

**Acknowledgements.** M. M. was partially supported by the Chilean Science Foundation *Fondecyt* under the Grant 1085162. S. R. and R. T. d A. thank the Swiss National Science Foundation for financial support.

## 2 The method of the weakly conjugate operator

In this section we recall the basic characteristics of the method of the weakly conjugate operator, as originally introduced and applied to partial differential operators in [7, 8]. We then present the abstract form of the extension developed in [26, 27]. The method works for unbounded operators, but for our purposes it is enough to assume  $H$  bounded.

### 2.1 The standard theory

We start by introducing some notations. The symbol  $\mathcal{H}$  stands for a Hilbert space with scalar product  $\langle \cdot, \cdot \rangle$  and norm  $\| \cdot \|$ . Given two Hilbert spaces  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , we denote by  $\mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$  the set of bounded operators from  $\mathcal{H}_1$  to  $\mathcal{H}_2$ , and put  $\mathcal{B}(\mathcal{H}) := \mathcal{B}(\mathcal{H}, \mathcal{H})$ . We assume that  $\mathcal{H}$  is endowed with a strongly continuous unitary group  $\{W_t\}_{t \in \mathbb{R}}$ . Its selfadjoint generator is denoted by  $A$  and has domain  $D(A)$ . In most of the applications  $A$  is unbounded.

**Definition 2.1.** *A bounded selfadjoint operator  $H$  in  $\mathcal{H}$  belongs to  $C^1(A; \mathcal{H})$  if one of the following equivalent condition is satisfied:*

- (i) the map  $\mathbb{R} \ni t \mapsto W_{-t}HW_t \in \mathcal{B}(\mathcal{H})$  is strongly differentiable,  
(ii) the sesquilinear form

$$D(A) \times D(A) \ni (f, g) \mapsto i \langle Hf, Ag \rangle - i \langle Af, Hg \rangle \in \mathbb{C}$$

is continuous when  $D(A)$  is endowed with the topology of  $\mathcal{H}$ .

We denote by  $B$  the strong derivative in (i), or equivalently the bounded selfadjoint operator associated with the extension of the form in (ii). The operator  $B$  provides a rigorous meaning to the commutator  $i[H, A]$ . We write  $B > 0$  if  $B$  is positive and injective, namely if  $\langle f, Bf \rangle > 0$  for all  $f \in \mathcal{H} \setminus \{0\}$ .

**Definition 2.2.** The operator  $A$  is weakly conjugate to the bounded selfadjoint operator  $H$  if  $H \in C^1(A; \mathcal{H})$  and  $B \equiv i[H, A] > 0$ .

For  $B > 0$  let us consider the completion  $\mathcal{B}$  of  $\mathcal{H}$  with respect to the norm  $\|f\|_{\mathcal{B}} := \langle f, Bf \rangle^{1/2}$ . The adjoint space  $\mathcal{B}^*$  of  $\mathcal{B}$  can be identified with the completion of  $B\mathcal{H}$  with respect to the norm  $\|g\|_{\mathcal{B}^*} := \langle g, B^{-1}g \rangle^{1/2}$ . One has then the continuous dense embeddings  $\mathcal{B}^* \hookrightarrow \mathcal{H} \hookrightarrow \mathcal{B}$ , and  $B$  extends to an isometric operator from  $\mathcal{B}$  to  $\mathcal{B}^*$ . Due to these embeddings it makes sense to assume that  $\{W_t\}_{t \in \mathbb{R}}$  restricts to a  $C_0$ -group in  $\mathcal{B}^*$ , or equivalently that it extends to a  $C_0$ -group in  $\mathcal{B}$ . Under this assumption (tacitly assumed in the sequel) we keep the same notation for these  $C_0$ -groups. The domain of the generator of the  $C_0$ -group in  $\mathcal{B}$  (resp.  $\mathcal{B}^*$ ) endowed with the graph norm is denoted by  $D(A, \mathcal{B})$  (resp.  $D(A, \mathcal{B}^*)$ ). In analogy with Definition 2.1 the requirement  $B \in C^1(A; \mathcal{B}, \mathcal{B}^*)$  means that the map  $\mathbb{R} \ni t \mapsto W_{-t}BW_t \in \mathcal{B}(\mathcal{B}, \mathcal{B}^*)$  is strongly differentiable, or equivalently that the sesquilinear form

$$D(A, \mathcal{B}) \times D(A, \mathcal{B}) \ni (f, g) \mapsto i \langle f, BAg \rangle - i \langle Af, Bg \rangle \in \mathbb{C}$$

is continuous when  $D(A, \mathcal{B})$  is endowed with the topology of  $\mathcal{B}$ . Here,  $\langle \cdot, \cdot \rangle$  denotes the duality between  $\mathcal{B}$  and  $\mathcal{B}^*$ . Finally let  $\mathcal{E}$  be the Banach space  $(D(A, \mathcal{B}^*), \mathcal{B}^*)_{1/2, 1}$  defined by real interpolation, see for example [1, Proposition 2.7.3]. One has then the natural continuous embeddings  $\mathcal{B}(\mathcal{H}) \subset \mathcal{B}(\mathcal{B}^*, \mathcal{B}) \subset \mathcal{B}(\mathcal{E}, \mathcal{E}^*)$  and the following results [8, Theorem 2.1]:

**Theorem 2.3.** Assume that  $A$  is weakly conjugate to  $H$  and that  $B \equiv i[H, A]$  belongs to  $C^1(A; \mathcal{B}, \mathcal{B}^*)$ . Then there exists a constant  $c > 0$  such that

$$|\langle f, (H - \lambda \mp i\mu)^{-1}f \rangle| \leq c \|f\|_{\mathcal{E}}^2$$

for all  $\lambda \in \mathbb{R}$ ,  $\mu > 0$  and  $f \in \mathcal{E}$ . In particular the spectrum of  $H$  is purely absolutely continuous.

The global limiting absorption principle plays an important role for partial differential operators, cf. [7, 13, 24, 30]. However, for the examples included sections 3 and 4, the space  $\mathcal{E}$  involved has a rather obscure meaning and cannot be greatly simplified, so we shall only state the spectral results.

## 2.2 The extension

The extension proposed in [26, 27] relies on the following observation. Assume that  $H$  is a bounded selfadjoint operator in a Hilbert space  $\mathcal{H}$ . Assume also that there exists a selfadjoint operator  $A$  such that  $H \in C^1(A; \mathcal{H})$ , and that

$$B \equiv i[H, A] = K^2 \tag{2.1}$$

for some selfadjoint operator  $K$ , which is bounded by hypothesis. It follows that  $i[H, A] \geq 0$  but injectivity may not be satisfied. So let us introduce the decomposition of the Hilbert space  $\mathcal{H} := \mathcal{K} \oplus \mathcal{G}$  with  $\mathcal{K} := \ker(K)$ . The operator  $B$  is reduced by this decomposition and its restriction  $B_0$  to  $\mathcal{G}$  satisfies  $B_0 > 0$ . Formally, the positivity and injectivity of  $B_0$  are rather promising, but are obviously not sufficient for a direct application of the MWCO to  $B_0$ .

However, let us already observe that some informations on  $\mathcal{H}_p(H)$  can be inferred from (2.1). Indeed, it follows from the Virial Theorem [1, Prop. 7.2.10] that any eigenvector  $f$  of  $H$  satisfies  $\langle f, i[H, A]f \rangle = 0$ . So  $0 = \langle f, K^2 f \rangle = \|Kf\|^2$ , i.e.  $f \in \ker(K)$ , and one has proved:

**Lemma 2.4.** *If  $H \in C^1(A; \mathcal{H})$  and  $i[H, A] = K^2$ , then  $\mathcal{H}_p(H) \subset \mathcal{K} = \ker(K)$ .*

Let us now come back to the analysis of  $B_0$ . The space  $\mathcal{B}$  can still be defined in analogy with what has been presented in the previous section:  $\mathcal{B}$  is the completion of  $\mathcal{G}$  with respect to the norm  $\|f\|_{\mathcal{B}} := \langle f, Bf \rangle^{1/2}$ . Then, the adjoint space  $\mathcal{B}^*$  of  $\mathcal{B}$  can be identified with the completion of  $B\mathcal{G}$  with respect to the norm  $\|g\|_{\mathcal{B}^*} := \langle g, B^{-1}g \rangle^{1/2}$ . But in order to go further on, some compatibility has to be imposed between the decomposition of the Hilbert space and the operators  $H$  and  $A$ . Let us assume that both operators  $H$  and  $A$  are reduced by the decomposition  $\mathcal{K} \oplus \mathcal{G}$  of  $\mathcal{H}$ , and let  $H_0$  and  $A_0$  denote their respective restriction to  $\mathcal{G}$ . It clearly follows from the above lemma that  $H_0$  has no point spectrum.

We are now in a suitable position to rephrase Theorem 2.3 in our present framework. We freely use the notations introduced above.

**Theorem 2.5.** *Assume that  $H \in C^1(A; \mathcal{H})$  and that  $B \equiv i[H, A] = K^2$ . Assume furthermore that both operators  $H$  and  $A$  are reduced by the decomposition  $\ker(K) \oplus \ker(K)^\perp$  of  $\mathcal{H}$  and that  $B_0$  belongs to  $C^1(A_0; \mathcal{B}, \mathcal{B}^*)$ . Then a limiting absorption principle holds for  $H_0$  uniformly on  $\mathbb{R}$ , and in particular the spectrum of  $H_0$  is purely absolutely continuous.*

A straightforward consequence of this statement is that

$$\mathcal{H}_{sc}(H) \subset \mathcal{H}_s(H) \subset \mathcal{K} = \ker(K).$$

We will see in the applications below that Theorem 2.5 applies to various situations, and that it really is a useful extension of the original method of the weakly conjugate operator.

### 3 Spectral analysis for adjacency operators on graphs

We present in this section the results of [26] on the spectral analysis for adjacency operators on graphs. We follow the notations and conventions of this paper regarding graph theory.

A *graph* is a couple  $(X, \sim)$  formed of a non-void countable set  $X$  and a symmetric relation  $\sim$  on  $X$  such that  $x \sim y$  implies  $x \neq y$ . The points  $x \in X$  are called vertices and couples  $(x, y) \in X \times X$  such that  $x \sim y$  are called edges. So, for simplicity, multiple edges and loops are forbidden in our definition of a graph. Occasionally  $(X, \sim)$  is said to be a simple graph. For any  $x \in X$  we denote by  $N(x) := \{y \in X : y \sim x\}$  the set of neighbours of  $x$ . We write  $\deg(x) := \#N(x)$  for the degree or valence of the vertex  $x$  and  $\deg(X) := \sup_{x \in X} \deg(x)$  for the degree of the graph. We also suppose that  $(X, \sim)$  is uniformly locally finite, *i.e.* that  $\deg(X) < \infty$ . A path from  $x$  to  $y$  is a sequence  $p = (x_0, x_1, \dots, x_n)$  of elements of  $X$ , usually denoted by  $x_0 x_1 \dots x_n$ , such that  $x_0 = x$ ,  $x_n = y$  and  $x_{j-1} \sim x_j$  for each  $j \in \{1, \dots, n\}$ .

Throughout this section we restrict ourselves to graphs  $(X, \sim)$  which are simple, infinite countable and uniformly locally finite. Given such a graph we consider the adjacency operator  $H$  acting in the Hilbert space  $\mathcal{H} := \ell^2(X)$  as

$$(Hf)(x) := \sum_{y \sim x} f(y), \quad f \in \mathcal{H}, \quad x \in X.$$

Due to [28, Theorem 3.1],  $H$  is a bounded selfadjoint operator with  $\|H\| \leq \deg(X)$  and spectrum  $\sigma(H) \subset [-\deg(X), \deg(X)]$ .

Results on the nature of the spectrum of adjacency operators on graphs are quite sparse. Some absolutely continuous examples are given in [28], including the lattice  $\mathbb{Z}^n$  and homogeneous trees. For cases in which singular components are present we refer to [11], [17], [31] and [32].

We now introduce the key concept of [26]. Sums over the empty set are zero by convention.

**Definition 3.1.** *A function  $\Phi : X \rightarrow \mathbb{R}$  is semi-adapted to the graph  $(X, \sim)$  if*

- (i) *there exists  $C \geq 0$  such that  $|\Phi(x) - \Phi(y)| \leq C$  for all  $x, y \in X$  with  $x \sim y$ ,*

(ii) for any  $x, y \in X$  one has

$$\sum_{z \in N(x) \cap N(y)} [2\Phi(z) - \Phi(x) - \Phi(y)] = 0. \quad (3.1)$$

If in addition for any  $x, y \in X$  one has

$$\sum_{z \in N(x) \cap N(y)} [\Phi(z) - \Phi(x)] [\Phi(z) - \Phi(y)] [2\Phi(z) - \Phi(x) - \Phi(y)] = 0, \quad (3.2)$$

then  $\Phi$  is adapted to the graph  $(X, \sim)$ .

For a function  $\Phi$  semi-adapted to the graph  $(X, \sim)$  we consider in  $\mathcal{H}$  the operator  $K$  given by

$$(Kf)(x) := i \sum_{y \sim x} [\Phi(y) - \Phi(x)] f(y), \quad f \in \mathcal{H}, x \in X.$$

The operator  $K$  is selfadjoint and bounded due to the condition (i) of Definition 3.1. It commutes with  $H$ , as a consequence of Condition (3.1). We also decompose the Hilbert space  $\mathcal{H}$  into the direct sum  $\mathcal{H} = \mathcal{K} \oplus \mathcal{G}$ , where  $\mathcal{G}$  is the closure of the range  $K\mathcal{H}$  of  $K$ , thus the orthogonal complement of the closed subspace

$$\mathcal{K} := \ker(K) = \left\{ f \in \mathcal{H} : \sum_{y \in N(x)} \Phi(y) f(y) = \Phi(x) \sum_{y \in N(x)} f(y) \text{ for each } x \in X \right\}.$$

It is shown in [26, Sec. 4] that  $H$  and  $K$  are reduced by this decomposition, and that their restrictions  $H_0$  and  $K_0$  to the Hilbert space  $\mathcal{G}$  are bounded selfadjoint operators.

A rather straightforward application of the general theory presented in 2.2 gives

**Theorem 3.2** (Theorem 3.2 of [26]). *Assume that  $\Phi$  is a function semi-adapted to the graph  $(X, \sim)$ . Then  $H_0$  has no point spectrum.*

**Theorem 3.3** (Theorem 3.3 of [26]). *Let  $\Phi$  be a function adapted to the graph  $(X, \sim)$ . Then the operator  $H_0$  has a purely absolutely continuous spectrum.*

The role of the weakly conjugate operator is played by  $A := \frac{1}{2}(\Phi K + K\Psi)$  and the assumptions imposed on  $\Phi$  make the general theory work.

For a certain class of admissible graphs, the result of Theorem 3.3 on the restriction  $H_0$  can be turned into a statement on the original adjacency operator  $H$ . The notion of admissibility requires (among other things) the graph to be directed. Thus the family of neighbors  $N(x) := \{y \in X : y \sim x\}$  is divided into two disjoint sets  $N^-(x)$  (fathers) and  $N^+(x)$  (sons),  $N(x) = N^-(x) \sqcup N^+(x)$ . We write  $y < x$  if  $y \in N^-(x)$  and  $x < y$  if  $y \in N^+(x)$ . On drawings, we set an arrow from  $y$  to  $x$  ( $x \leftarrow y$ ) if  $x < y$ , and say that the edge from  $y$  to  $x$  is positively oriented.

We assume that the directed graph subjacent to  $X$ , from now on denoted by  $(X, <)$ , is *admissible* with respect to these decompositions, *i.e.* (i) it admits a position function and (ii) it is uniform. A *position function* is a function  $\Phi : X \rightarrow \mathbb{Z}$  such that  $\Phi(y) + 1 = \Phi(x)$  whenever  $y < x$ . It is easy to see that it exists if and only if all paths between two points have the same index (which is the difference between the number of positively and negatively oriented edges). The directed graph  $(X, <)$  is called *uniform* if for any  $x, y \in X$  the number  $\# [N^-(x) \cap N^-(y)]$  of common fathers of  $x$  and  $y$  equals the number  $\# [N^+(x) \cap N^+(y)]$  of common sons of  $x$  and  $y$ . Thus the admissibility of a directed graph is an explicit property that can be checked directly, without making any choice. The graph  $(X, \sim)$  is admissible if there exists an admissible directed graph subjacent to it.

**Theorem 3.4** (Theorem 1.1 of [26]). *The adjacency operator of an admissible graph  $(X, \sim)$  is purely absolutely continuous, except at the origin, where it may have an eigenvalue with eigenspace*

$$\ker(H) = \left\{ f \in \mathcal{H} : \sum_{y < x} f(y) = 0 = \sum_{y > x} f(y) \text{ for each } x \in X \right\}. \quad (3.3)$$

Many examples of periodic graphs, both admissible and non-admissible, are presented in [26, Sec. 6]. In particular, it is explained that periodicity does not lead automatically to absolute continuity, especially (but not only) if the number of orbits is infinite.  $D$ -products of graphs, as well as the graph associated with the one-dimensional XY Hamiltonian, are also treated in [26]. We recall in Figures 1, 2, and 5 some two-dimensional  $\mathbb{Z}$ -periodic examples taken from [26, Sec. 6]. More involved,  $\mathbb{Z}^n$ -periodic situations are also available.




Figure 1: Example of an admissible directed graph with  $\ker(H) \neq \{0\}$




Figure 2: Example of an admissible directed graph




Figure 3: Examples of admissible, directed graphs with  $\ker(H) = \{0\}$




Figure 4: Example of an admissible, directed graph with  $\ker(H) \neq \{0\}$




Figure 5: Example of a non-admissible, adapted graph (with function  $\Phi$  as indicated)

## 4 Convolution operators on locally compact groups

In this section we consider locally compact groups  $X$ , abelian or not, and convolution operators  $H_\mu$ , acting on  $L^2(X)$ , defined by suitable measures  $\mu$  belonging to  $M(X)$ , the Banach  $*$ -algebra of complex bounded Radon measures on  $X$ . Using the method of the weakly conjugate operator, we determine subspaces  $\mathcal{K}_\mu^1$  and  $\mathcal{K}_\mu^2$  of  $L^2(X)$ , explicitly defined in terms of  $\mu$  and the family  $\text{Hom}(X, \mathbb{R})$  of continuous group morphisms  $\Phi : X \rightarrow \mathbb{R}$ , such that  $\mathcal{H}_p(H_\mu) \subset \mathcal{K}_\mu^1$  and  $\mathcal{H}_s(H_\mu) \subset \mathcal{K}_\mu^2$ . This result, obtained in [27], supplements other works on the spectral theory of operators on groups and graphs [4, 5, 6, 9, 10, 11, 15, 17, 18, 19, 21, 22, 23, 31].

Let  $X$  be a locally compact group (LCG) with identity  $e$ , center  $Z(X)$  and modular function  $\Delta$ . Let us fix a left Haar measure  $\lambda$  on  $X$ , using the notation  $dx := d\lambda(x)$ . On discrete groups the counting measure (assigning mass 1 to every point) is considered. The notation *a.e.* stands for “almost everywhere” and refers to the Haar measure  $\lambda$ .

We consider in the sequel the convolution operator  $H_\mu$ ,  $\mu \in M(X)$ , acting in the Hilbert space  $\mathcal{H} := L^2(X, d\lambda)$ , *i.e.*

$$H_\mu f := \mu * f,$$

where  $f \in \mathcal{H}$  and

$$(\mu * f)(x) := \int_X d\mu(y) f(y^{-1}x) \quad \text{for a.e. } x \in X.$$

The operator  $H_\mu$  is bounded with norm  $\|H_\mu\| \leq \|\mu\|$ , and it admits an adjoint operator  $H_\mu^*$  equal to  $H_{\mu^*}$ , the convolution operator by  $\mu^* \in M(X)$  defined by  $\mu^*(E) = \overline{\mu(E^{-1})}$ . If the measure  $\mu$  is absolutely continuous w.r.t. the left Haar measure  $\lambda$ , so that  $d\mu = a d\lambda$  with  $a \in L^1(X)$ , then  $\mu^*$  is also absolutely continuous w.r.t.  $\lambda$  and  $d\mu^* = a^* d\lambda$ , where  $a^*(x) := \overline{a(x^{-1})} \Delta(x^{-1})$  for *a.e.*  $x \in X$ . In such a case we simply write  $H_a$  for  $H_{a d\lambda}$ . We shall always assume that  $H_\mu$  is selfadjoint, that is  $\mu = \mu^*$ .

Given  $\mu \in M(X)$ , let  $\varphi : X \rightarrow \mathbb{R}$  be such that the linear functional

$$F : C_0(X) \rightarrow \mathbb{C}, \quad g \mapsto \int_X d\mu(x) \varphi(x) g(x)$$

is bounded. Then there exists a unique measure in  $M(X)$  associated to  $F$ , due to the Riesz-Markov representation theorem. We write  $\varphi\mu$  for this measure, and we simply say that  $\varphi$  is such that  $\varphi\mu \in M(X)$ . We call *real character* any continuous group morphism  $\Phi : X \rightarrow \mathbb{R}$ .

**Definition 4.1.** Let  $\mu = \mu^* \in M(X)$ .

(a) A real character  $\Phi$  is semi-adapted to  $\mu$  if  $\Phi\mu, \Phi^2\mu \in M(X)$ , and  $(\Phi\mu) * \mu = \mu * (\Phi\mu)$ . The set of real characters that are semi-adapted to  $\mu$  is denoted by  $\text{Hom}_\mu^1(X, \mathbb{R})$ .

(b) A real character  $\Phi$  is adapted to  $\mu$  if  $\Phi$  is semi-adapted to  $\mu$ ,  $\Phi^3\mu \in M(X)$ , and  $(\Phi\mu) * (\Phi^2\mu) = (\Phi^2\mu) * (\Phi\mu)$ . The set of real characters that are adapted to  $\mu$  is denoted by  $\text{Hom}_\mu^2(X, \mathbb{R})$ .

Let  $\mathcal{K}_\mu^j := \bigcap_{\Phi \in \text{Hom}_\mu^j(X, \mathbb{R})} \ker(H_{\Phi\mu})$ , for  $j = 1, 2$ ; then the main result is the following.

**Theorem 4.2** (Theorem 2.2 of [27]). Let  $X$  be a LCG and let  $\mu = \mu^* \in M(X)$ . Then

$$\mathcal{H}_p(H_\mu) \subset \mathcal{K}_\mu^1 \quad \text{and} \quad \mathcal{H}_s(H_\mu) \subset \mathcal{K}_\mu^2.$$

The cases  $\mathcal{K}_\mu^1 = \{0\}$  or  $\mathcal{K}_\mu^2 = \{0\}$  are interesting; in the first case  $H_\mu$  has no eigenvalues, and in the second case  $H_\mu$  is purely absolutely continuous. A more precise result is obtained in a particular situation.

**Corollary 4.3** (Corollary 2.3 of [27]). Let  $X$  be a LCG and let  $\mu = \mu^* \in M(X)$ . Assume that there exists a real character  $\Phi$  adapted to  $\mu$  such that  $\Phi^2$  is equal to a nonzero constant on  $\text{supp}(\mu)$ . Then  $H_\mu$  has a purely absolutely continuous spectrum, with the possible exception of an eigenvalue located at the origin, with eigenspace  $\ker(H_\mu) = \ker(H_{\Phi\mu})$ .



Corollary 4.3 specially applies to adjacency operators on certain classes of Cayley graphs, which are Hecke-type operators in the regular representation, thus convolution operators on discrete groups.

To see how the method of the weakly conjugate operator comes into play, let us sketch the proof of the second inclusion in Theorem 4.2. In quantum mechanics in  $\mathbb{R}^d$ , the position operators  $Q_j$  and the momentum operators  $P_j$  satisfy the relation  $P_j = i[H, Q_j]$  with  $H := -\frac{1}{2}\Delta$ , and the usual conjugate operator is the generator of dilations  $D := \frac{1}{2}\sum_j(Q_j P_j + P_j Q_j)$ . So if we regard  $\Phi \in \text{Hom}_\mu^2(X, \mathbb{R})$  as a position operator on  $X$ , it is reasonable to think of  $K := i[H_\mu, \Phi]$  as the corresponding momentum operator and to use  $A := \frac{1}{2}(\Phi K + K\Phi)$  as a tentative conjugate operator. In fact, simple calculations using the hypotheses of Definition 4.1.(b) show that

$$K = -iH_{\Phi\mu} \in \mathcal{B}(\mathcal{H}) \quad \text{and} \quad i[H_\mu, A] = K^2.$$

Therefore  $H_\mu \in C^1(A; \mathcal{H})$  and the commutator  $i[H_\mu, A]$  is a positive operator. However, in order to apply the theory of Section 2, we need strict positivity. So, we consider the subspace  $\mathcal{G} := [\ker(K)]^\perp$  of  $\mathcal{H}$ , where  $i[H_\mu, A] \equiv K^2$  is strictly positive. The orthogonal decomposition  $\mathcal{H} := \mathcal{K} \oplus \mathcal{G}$  with  $\mathcal{K} := \ker(K)$ , reduces  $H_\mu$ ,  $K$ , and  $A$ , and their restrictions  $H_0$ ,  $K_0$ , and  $A_0$  to  $\mathcal{G}$  are selfadjoint. It turns out that all the other conditions necessary to apply Theorem 2.5 can also be verified in the Hilbert space  $\mathcal{G}$ . So we get the inclusion

$$\mathcal{H}_s \subset \mathcal{K} = \ker(H_{\Phi\mu}).$$

Since  $\Phi \in \text{Hom}_\mu^2(X, \mathbb{R})$  is arbitrary, this implies the second inclusion of Theorem 4.2.

## 4.1 Examples

The construction of weakly conjugate operators for  $H_\mu$  relies on real characters. So a small vector space  $\text{Hom}(X, \mathbb{R})$  is an obstacle to applying the method. A real character  $\Phi$  maps compact subgroups of  $X$  to the unique compact subgroup  $\{0\}$  of  $\mathbb{R}$ . Consequently, abundance of compact elements (elements  $x \in X$  generating compact subgroups) prevents us from constructing weakly conjugate operators. The extreme case is when  $X$  is itself compact, so that  $\text{Hom}(X, \mathbb{R}) = \{0\}$ . Actually in such a case all convolution operators  $H_\mu$  have pure point spectrum. We review now briefly some of the groups for which we succeeded in [27] to apply the MWCO.

If the group  $X$  is unimodular, one can exploit commutativity in a non-commutative setting by using central measures (*i.e.* elements of the center  $Z[M(X)]$  of the convolution Banach \*-algebra  $M(X)$ ). For instance, in the case of central groups [16], we have the following result ( $\mathcal{B}(X)$  stands for the closed subgroup generated by the set of compact elements of  $X$ ):

**Proposition 4.4** (Proposition 4.2 of [27]). *Let  $X$  be a central group and  $\mu_0 = \mu_0^* \in M(X)$  a central measure such that  $\text{supp}(\mu_0)$  is compact and not included in  $\mathcal{B}(X)$ . Let  $\mu_1 = \mu_1^* \in M(X)$  with  $\text{supp}(\mu_1) \subset \mathcal{B}(X)$  and set  $\mu := \mu_0 + \mu_1$ . Then  $\mathcal{H}_{\text{ac}}(H_\mu) \neq \{0\}$ .*

Let us recall three examples deduced from Proposition 4.4 taken from [27].

**Example 4.5.** *Let  $X := S_3 \times \mathbb{Z}$ , where  $S_3$  is the symmetric group of degree 3. The group  $S_3$  has a presentation  $\langle a, b \mid a^2, b^2, (ab)^3 \rangle$ , and its conjugacy classes are  $E_1 = E_1^{-1} = \{e\}$ ,  $E_2 = E_2^{-1} = \{a, b, aba\}$  and  $E_3 = E_3^{-1} = \{ab, ba\}$ . Set  $\mathcal{E} := \{E_2, E_3\}$  and choose  $I_{E_1}, I_{E_2}$  two finite symmetric subsets of  $\mathbb{Z}$ , each of them containing at least two elements. Then  $\mathcal{H}_{\text{ac}}(H_{\chi_S}) \neq \{0\}$  if  $S := \bigcup_{E \in \mathcal{E}} E \times I_E$ .*

**Example 4.6.** *Let  $X := \text{SU}(2) \times \mathbb{R}$ , where  $\text{SU}(2)$  is the group (with Haar measure  $\lambda_2$ ) of  $2 \times 2$  unitary matrices of determinant +1. For each  $\vartheta \in [0, \pi]$  let  $C(\vartheta)$  be the conjugacy class of the matrix  $\text{diag}(e^{i\vartheta}, e^{-i\vartheta})$  in  $\text{SU}(2)$ . A direct calculation (using for instance Euler angles) shows that  $\lambda_2(\bigcup_{\vartheta \in J} C(\vartheta)) > 0$  for each  $J \subset [0, \pi]$  with nonzero Lebesgue measure. Set  $E_1 := \bigcup_{\vartheta \in (0,1)} C(\vartheta)$ ,  $E_2 := \bigcup_{\vartheta \in (2,\pi)} C(\vartheta)$ ,  $\mathcal{E} := \{E_1, E_2\}$ ,  $I_{E_1} := (-1, 1)$ , and  $I_{E_2} := (-3, -2) \cup (2, 3)$ . Then  $\mathcal{H}_{\text{ac}}(H_{\chi_S}) \neq \{0\}$  if  $S := \bigcup_{E \in \mathcal{E}} E \times I_E$ .*

**Example 4.7.** *Let  $X$  be a central group, let  $z \in Z(X) \setminus \mathcal{B}(X)$ , and set  $\mu := \delta_z + \delta_{z^{-1}} + \mu_1$  for some  $\mu_1 = \mu_1^* \in M(X)$  with  $\text{supp}(\mu_1) \subset \mathcal{B}(X)$ . Then  $\mu$  satisfies the hypotheses of Proposition 4.4, and we can choose  $\Phi \in \text{Hom}(X, \mathbb{R})$  such that  $\Phi(z) = \frac{1}{2}\Phi(z^2) \neq 0$  (note in particular that  $z \notin \mathcal{B}(X)$  iff  $z^2 \notin \mathcal{B}(X)$  and that  $\Phi\mu_1 = 0$ ). Thus  $\mathcal{H}_s(H_\mu) \subset \ker(H_{\Phi\mu})$ . But  $f \in \mathcal{H}$  belongs to  $\ker(H_{\Phi\mu}) = \ker(H_{\Phi(\delta_z + \delta_{z^{-1}})})$  iff*



$f(z^{-1}x) = f(zx)$  for a.e.  $x \in X$ . This periodicity w.r.t. the non-compact element  $z^2$  implies that the  $L^2$ -function  $f$  should vanish a.e. and thus that  $\mathcal{H}_{\text{ac}}(H_\mu) = \mathcal{H}$ .

If  $X$  is abelian all the commutation relations in Definition 4.1 are satisfied. Moreover one can use the Fourier transform  $\mathcal{F}$  to map unitarily  $H_\mu$  on the operator  $M_m$  of multiplication with  $m = \mathcal{F}(\mu)$  on the dual group  $\widehat{X}$  of  $X$ . So one gets from Theorem 4.2 a general lemma on multiplication operators. We recall some definitions before stating it.

**Definition 4.8.** *The function  $m : \widehat{X} \rightarrow \mathbb{C}$  is differentiable at  $\xi \in \widehat{X}$  along the one-parameter subgroup  $\varphi \in \text{Hom}(\mathbb{R}, \widehat{X})$  if the function  $\mathbb{R} \ni t \mapsto m(\xi + \varphi(t)) \in \mathbb{C}$  is differentiable at  $t = 0$ . In such a case we write  $(d_\varphi m)(\xi)$  for  $\frac{d}{dt} m(\xi + \varphi(t))|_{t=0}$ . Higher order derivatives, when existing, are denoted by  $d_\varphi^k m$ ,  $k \in \mathbb{N}$ .*

We say that the one-parameter subgroup  $\varphi : \mathbb{R} \rightarrow \widehat{X}$  is in  $\text{Hom}_{m_0}^1(\mathbb{R}, \widehat{X})$  if  $m$  is twice differentiable w.r.t.  $\varphi$  and  $d_\varphi m, d_\varphi^2 m \in \mathcal{F}(M(X))$ . If, in addition,  $m$  is three times differentiable w.r.t.  $\varphi$  and  $d_\varphi^3 m \in \mathcal{F}(M(X))$  too, we say that  $\varphi$  belongs to  $\text{Hom}_{m_0}^2(\mathbb{R}, \widehat{X})$ .

**Lemma 4.9** (Corollary 4.7 of [27]). *Let  $X$  be a locally compact abelian group and let  $m_0, m_1$  be real functions with  $\mathcal{F}^{-1}(m_0), \mathcal{F}^{-1}(m_1) \in M(X)$  and  $\text{supp}(\mathcal{F}^{-1}(m_1)) \subset \mathcal{B}(X)$ . Then*

$$\mathcal{H}_p(M_{m_0+m_1}) \subset \bigcap_{\varphi \in \text{Hom}_{m_0}^1(\mathbb{R}, \widehat{X})} \ker(M_{d_\varphi m_0})$$

and

$$\mathcal{H}_s(M_{m_0+m_1}) \subset \bigcap_{\varphi \in \text{Hom}_{m_0}^2(\mathbb{R}, \widehat{X})} \ker(M_{d_\varphi m_0}).$$

We end up this section by considering a class of semidirect products. Let  $N, G$  be two discrete groups with  $G$  abelian (for which we use additive notations), and let  $\tau : G \rightarrow \text{Aut}(N)$  be a group morphism. Let  $X := N \times_\tau G$  be the  $\tau$ -semidirect product of  $N$  by  $G$ . The multiplication in  $X$  is defined by

$$(n, g)(m, h) := (n\tau_g(m), g + h),$$

so that

$$(n, g)^{-1} = (\tau_{-g}(n^{-1}), -g).$$

In this situation it is shown in [27] that many convolution operators  $H_a$ , with  $a : X \rightarrow \mathbb{C}$  of finite support, have a non-trivial absolutely continuous component. For instance, we have the following for a type of wreath products.

**Example 4.10.** *Take  $G$  a discrete abelian group and put  $N := R^J$ , where  $R$  is an arbitrary discrete group and  $J$  is a finite set on which  $G$  acts by  $(g, j) \mapsto g(j)$ . Then  $\tau_g(\{r_j\}_{j \in J}) := \{r_{g(j)}\}_{j \in J}$  defines an action of  $G$  on  $R^J$ , thus we can construct the semidirect product  $R^J \times_\tau G$ . If  $G_0 = -G_0 \subset G$  and  $R_0 = R_0^{-1} \subset R$  are finite subsets with  $G_0 \cap [G \setminus \mathcal{B}(G)] \neq \emptyset$ , then  $N_0 := R_0^{G_0}$  satisfies all the conditions of [27, Sec. 4.4]. Thus  $\mathcal{H}_{\text{ac}}(H_{\chi_S}) \neq \{0\}$  if  $S := N_0 \times G_0$ .*

Virtually the methods of [27] could also be applied to non-split group extensions.

## References

- [1] W. O. Amrein, A. Boutet de Monvel and V. Georgescu.  *$C_0$ -groups, commutator methods and spectral theory of  $N$ -body Hamiltonians*. Volume 135 of *Progress in Math.*, Birkhäuser, Basel, 1996.
- [2] M. A. Astaburuaga, O. Bourget, V. Cortés and C. Fernández. Absence of point spectrum for unitary operators. *J. Differential Equations* 244(2): 229–241, 2008.

- [3] M. A. Astaburuaga, O. Bourget, V. Cortés and C. Fernández. Floquet operators without singular continuous spectrum. *J. Funct. Anal.* 238(2): 489–517, 2006.
- [4] L. Bartholdi and R. I. Grigorchuk. On the spectrum of Hecke type operators related to some fractal groups. *Tr. Mat. Inst. Steklova* 231 (Din. Sist., Avtom. i Beskon. Gruppy): 5–45, 2000.
- [5] L. Bartholdi and W. Woess. Spectral computations on lamplighter groups and Diestel-Leader graphs. *J. Fourier Anal. Appl.* 11(2): 175–202, 2005.
- [6] C. Béguin, A. Valette and A. Zuk. On the spectrum of a random walk on the discrete Heisenberg group and the norm of Harper’s operator. *J. Geom. Phys.* 21(4): 337–356, 1997.
- [7] A. Boutet de Monvel, G. Kazantseva and M. Măntoiu. Some anisotropic Schrödinger operators without singular spectrum. *Helv. Phys. Acta* 69(1): 13–25, 1996.
- [8] A. Boutet de Monvel and M. Măntoiu. The method of the weakly conjugate operator. In *Inverse and algebraic quantum scattering theory (Lake Balaton, 1996)*, volume 488 of *Lecture Notes in Phys.*, pages 204–226, Springer, Berlin, 1997.
- [9] J. Breuer. Singular continuous spectrum for the Laplacian on a certain sparse tree. *Commun. Math. Phys.* 269: 851–857, 2007.
- [10] J. Breuer. Singular continuous and dense point spectrum for sparse tree with finite dimensions. In *Probability and Mathematical Physics* (eds. D. Dawson, V. Jaksic and B. Vainberg), CRM Proc. and Lecture Notes 42, pages 65–83, 2007.
- [11] W. Dicks and T. Schick. The spectral measure of certain elements of the complex group ring of a wreath product. *Geom. Dedicata* 93: 121–137, 2002.
- [12] S. Fournais and E. Skibsted. Zero energy asymptotics of the resolvent for a class of slowly decaying potentials. *Math. Z.* 248: 593–633, 2004.
- [13] A. Iftimovici and M. Măntoiu. Limiting absorption principle at critical values for the Dirac operator. *Lett. Math. Phys.* 49(3): 235–243, 1999.
- [14] V. Georgescu, C. Gérard and J.S. Moeller. Commutators,  $C_0$ -semigroups and Resolvent Estimates. *J. Funct. Anal.* 216(2): 303–361, 2004.
- [15] G. Georgescu and S. Golénia. Isometries, Fock spaces, and spectral analysis of Schrödinger operators on trees. *J. Funct. Anal.* 227(2): 389–429, 2005.
- [16] S. Grosser and M. Moskowitz. On central topological groups. *Trans. Amer. Math. Soc.* 127: 317–340, 1967.
- [17] R. I. Grigorchuk and A. Žuk. The lamplighter group as a group generated by a 2-state automaton, and its spectrum. *Geom. Dedicata* 87(1-3): 209–244, 2001.
- [18] P. de la Harpe, A. Gyan Robertson and A. Valette. On the spectrum of the sum of generators for a finitely generated group. *Israel J. Math.* 81(1-2): 65–96, 1993.
- [19] P. de la Harpe, A. Gyan Robertson and A. Valette. On the spectrum of the sum of generators of a finitely generated group. II. *Colloq. Math.* 65(1): 87–102, 1993.
- [20] I. Herbst. Spectral and scattering theory for Schrödinger operators with potentials independent of  $|x|$ . *Amer. J. Math.* 113: 509–565, 1991.
- [21] M. Kambites, P. V. Silva and B. Steinberg. The spectra of lamplighter groups and Cayley machines. *Geom. Dedicata* 120: 193–227, 2006.

- [22] H. Kesten. Symmetric random walks on groups. *Trans. Amer. Math. Soc.* 92: 336–354, 1959.
- [23] L. Malozemov and A. Teplyaev. Pure point spectrum of the Laplacians on fractal graphs. *J. Funct. Anal.* 129: 390–405, 1995.
- [24] M. Măntoiu and M. Pascu. Global resolvent estimates for multiplication operators. *J. Operator. Th.* 36: 283–294, 1996.
- [25] M. Măntoiu and S. Richard. Absence of singular spectrum for Schrödinger operators with anisotropic potentials and magnetic fields. *J. Math. Phys.* 41(5): 2732–2740, 2000.
- [26] M. Măntoiu, S. Richard and R. Tiedra de Aldecoa. Spectral analysis for adjacency operators on graphs. *Ann. Henri Poincaré* 8(7): 1401–1423, 2007.
- [27] M. Măntoiu and R. Tiedra de Aldecoa. Spectral analysis for convolution operators on locally compact groups. *J. Funct. Anal.* 253(2): 675–691, 2007.
- [28] B. Mohar and W. Woess. A survey on spectra of infinite graphs. *Bull. London Math. Soc.* 21(3): 209–234, 1989.
- [29] M. Reed and B. Simon. *Methods of modern mathematical physics, IV: Analysis of Operators*. Academic Press, New York, 1978.
- [30] S. Richard. Some improvements in the method of the weakly conjugate operator. *Lett. Math. Phys.* 76: 27–36, 2006.
- [31] B. Simon. Operators with singular continuous spectrum VI: Graph Laplacians and Laplace-Beltrami operators. *Proc. Amer. Math. Soc.* 124: 1177–1182, 1996.
- [32] I. Veselić. Spectral analysis of percolation hamiltonians. *Math. Ann* 331: 841–865, 2005.