

$P=W$

Tamás Hausel

Institute of Science and Technology Austria  
<http://hausel.ist.ac.at>

Between Topology and Quantum Field Theory  
A conference in celebration of Dan Freed's 60th birthday  
Austin  
January 2019



- $X$  smooth projective complex algebraic variety

# Hodge theory

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology

# Hodge theory

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$
- $H_B^1(X; \mathbb{C}) \cong \text{Hom}(H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}) \cong \text{Hom}(\pi_1(X) \rightarrow \mathbb{C})$

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$
- $H_B^1(X; \mathbb{C}) \cong \text{Hom}(H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}) \cong \text{Hom}(\pi_1(X) \rightarrow \mathbb{C})$
- $H_{\text{Dol}}^1(X; \mathbb{C}) \cong H^1(X; \mathcal{O}_X) \oplus H^0(X; \Omega_X^1)$



- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$
- $H_B^1(X; \mathbb{C}) \cong \text{Hom}(H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}) \cong \text{Hom}(\pi_1(X) \rightarrow \mathbb{C})$
- $H_{\text{Dol}}^1(X; \mathbb{C}) \cong H^1(X; \mathcal{O}_X) \oplus H^0(X; \Omega_X^1)$
- Hodge theory also implies

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$
- $H_B^1(X; \mathbb{C}) \cong \text{Hom}(H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}) \cong \text{Hom}(\pi_1(X) \rightarrow \mathbb{C})$
- $H_{\text{Dol}}^1(X; \mathbb{C}) \cong H^1(X; \mathcal{O}_X) \oplus H^0(X; \Omega_X^1)$
- Hodge theory also implies *Hard Lefschetz theorem*

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$
- $H_B^1(X; \mathbb{C}) \cong \text{Hom}(H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}) \cong \text{Hom}(\pi_1(X) \rightarrow \mathbb{C})$
- $H_{\text{Dol}}^1(X; \mathbb{C}) \cong H^1(X; \mathcal{O}_X) \oplus H^0(X; \Omega_X^1)$
- Hodge theory also implies *Hard Lefschetz theorem*

$$L^l : H^{d_X-l}(X; \mathbb{C}) \xrightarrow{\cong} H^{d_X+l}(X; \mathbb{C})$$

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$
- $H_B^1(X; \mathbb{C}) \cong \text{Hom}(H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}) \cong \text{Hom}(\pi_1(X) \rightarrow \mathbb{C})$
- $H_{\text{Dol}}^1(X; \mathbb{C}) \cong H^1(X; \mathcal{O}_X) \oplus H^0(X; \Omega_X^1)$
- Hodge theory also implies *Hard Lefschetz theorem*

$$L^l : \begin{array}{ccc} H^{d_X-l}(X; \mathbb{C}) & \xrightarrow{\cong} & H^{d_X+l}(X; \mathbb{C}) \\ x & \mapsto & x \cup \alpha^l \end{array}$$

- $X$  smooth projective complex algebraic variety
- $H_B^k(X; \mathbb{C})$  singular, or Betti cohomology
- $H_{\text{Dol}}^k(X; \mathbb{C}) = \bigoplus_{p+q=k} H^p(X; \Omega_X^q)$  Dolbeault cohomology
- Hodge theory implies  $H_B^k(X; \mathbb{C}) \cong H_{\text{Dol}}^k(X; \mathbb{C})$
- $H_B^1(X; \mathbb{C}) \cong \text{Hom}(H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}) \cong \text{Hom}(\pi_1(X) \rightarrow \mathbb{C})$
- $H_{\text{Dol}}^1(X; \mathbb{C}) \cong H^1(X; \mathcal{O}_X) \oplus H^0(X; \Omega_X^1)$
- Hodge theory also implies *Hard Lefschetz theorem*

$$L^l : \begin{array}{ccc} H^{d_X-l}(X; \mathbb{C}) & \xrightarrow{\cong} & H^{d_X+l}(X; \mathbb{C}) \\ X & \mapsto & X \cup \alpha^l \end{array}$$

for  $\alpha \in H^2(X; \mathbb{C})$  an ample class



- $C$  genus  $g$  curve

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$



- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\mathcal{M}_B^n$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\mathcal{M}_B^n := H_B^1(C; \mathrm{GL}_n)^d$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\mathcal{M}_B^n := H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned}\mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n\end{aligned}$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned}\mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n\end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned}\mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n\end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \end{array} \right.$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \begin{cases} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles } (E, \phi) \end{cases}$$



- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles } (E, \phi) \\ \phi \in H^0(C, \mathrm{End}(E) \otimes \Omega_C^1) \text{ Higgs field} \end{array} \right\}$$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles } (E, \phi) \\ \phi \in H^0(C, \mathrm{End}(E) \otimes \Omega_C^1) \text{ Higgs field} \end{array} \right\}$$

when  $(d, n) = 1$  these are smooth non-compact varieties

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles } (E, \phi) \\ \phi \in H^0(C, \mathrm{End}(E) \otimes \Omega_C^1) \text{ Higgs field} \end{array} \right\}$$

when  $(d, n) = 1$  these are smooth non-compact varieties

- Non-Abelian Hodge Theorem:  $\mathcal{M}_{\mathrm{Dol}}^n \stackrel{\mathrm{diff}}{\cong} \mathcal{M}_B^n$

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles } (E, \phi) \\ \phi \in H^0(C, \mathrm{End}(E) \otimes \Omega_C^1) \text{ Higgs field} \end{array} \right\}$$

when  $(d, n) = 1$  these are smooth non-compact varieties

- Non-Abelian Hodge Theorem:  $\mathcal{M}_{\mathrm{Dol}}^n \stackrel{\mathrm{diff}}{\cong} \mathcal{M}_B^n$   
(Hitchin, Donaldson, Corlette, Simpson)

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles } (E, \phi) \\ \phi \in H^0(C, \mathrm{End}(E) \otimes \Omega_C^1) \text{ Higgs field} \end{array} \right\}$$

when  $(d, n) = 1$  these are smooth non-compact varieties

- Non-Abelian Hodge Theorem:  $\mathcal{M}_{\mathrm{Dol}}^n \stackrel{\mathrm{diff}}{\cong} \mathcal{M}_B^n$   
(Hitchin, Donaldson, Corlette, Simpson)
- Problem

- $C$  genus  $g$  curve; fix  $n \in \mathbb{Z}_{>0}$  and  $d \in \mathbb{Z}$

$$\begin{aligned} \mathcal{M}_B^n &:= H_B^1(C; \mathrm{GL}_n)^d = [\mathrm{Hom}(\pi_1(C) \rightarrow \mathrm{GL}_n)]^d = \\ &= \{A_1, B_1, \dots, A_g, B_g \in \mathrm{GL}_n \mid \prod_{i=1}^g A_i^{-1} B_i^{-1} A_i B_i = e^{\frac{2\pi i d}{n}} \mathrm{Id}\} // \mathrm{PGL}_n \end{aligned}$$

$$\mathcal{M}_{\mathrm{Dol}}^n := H_{\mathrm{Dol}}^1(C; \mathrm{GL}_n)^d = \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles } (E, \phi) \\ \phi \in H^0(C, \mathrm{End}(E) \otimes \Omega_C^1) \text{ Higgs field} \end{array} \right\}$$

when  $(d, n) = 1$  these are smooth non-compact varieties

- Non-Abelian Hodge Theorem:  $\mathcal{M}_{\mathrm{Dol}}^n \stackrel{\text{diff}}{\cong} \mathcal{M}_B^n$   
(Hitchin, Donaldson, Corlette, Simpson)
- Problem: understand  $H^*(\mathcal{M}_B^n; \mathbb{C}) \cong H^*(\mathcal{M}_{\mathrm{Dol}}^n; \mathbb{C})$



- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$



- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$

# Weight filtration

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*
- $E(X; q) = q^{d_X} WH(1/q, -1)$

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*
- $E(X; q) = q^{d_X} WH(1/q, -1)$  *E-polynomial of  $X$*

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*
- $E(X; q) = q^{d_X} WH(1/q, -1)$  *E-polynomial of  $X$  (for smooth  $X$ )*



- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*
- $E(X; q) = q^{d_X} WH(1/q, -1)$  *E-polynomial of  $X$  (for smooth  $X$ )*

Theorem (Katz 2008)

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*
- $E(X; q) = q^{d_X} WH(1/q, -1)$  *E-polynomial of  $X$  (for smooth  $X$ )*

## Theorem (Katz 2008)

*If  $M$  is a smooth quasi-projective variety defined over  $\mathbb{Z}$*

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X)))t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*
- $E(X; q) = q^{d_X} WH(1/q, -1)$  *E-polynomial of  $X$  (for smooth  $X$ )*

## Theorem (Katz 2008)

*If  $M$  is a smooth quasi-projective variety defined over  $\mathbb{Z}$  and*

$$\#\{M(\mathbb{F}_q)\} = E(q)$$

*is a polynomial in  $q$ ,*

- (Deligne 1971) proved the existence of  $W_0 \subset \cdots \subset W_i \subset \cdots \subset W_{2k} = H^k(X; \mathbb{Q})$  for any complex algebraic variety  $X$
- $WH(X; q, t) = \sum \dim(W_i/W_{i-1}(H^k(X))) t^k q^{\frac{i}{2}}$  *weight polynomial*
- $P(X; t) = WH(X; 1, t)$  *Poincaré polynomial*
- $E(X; q) = q^{d_X} WH(1/q, -1)$  *E-polynomial of  $X$  (for smooth  $X$ )*

## Theorem (Katz 2008)

*If  $M$  is a smooth quasi-projective variety defined over  $\mathbb{Z}$  and*

$$\#\{M(\mathbb{F}_q)\} = E(q)$$

*is a polynomial in  $q$ , then  $E(M; q) = E(q)$ .*

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

*Let  $z \in G$  in a finite group  $G$*

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

*Let  $z \in G$  in a finite group  $G$  then*

$$\frac{1}{|G|} \#\{a_j, b_j \in G \mid [a_1, b_1] \dots [a_g, b_g] = z\}$$

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

Let  $z \in G$  in a finite group  $G$  then

$$\begin{aligned} \frac{1}{|G|} \#\{a_j, b_j \in G \mid [a_1, b_1] \dots [a_g, b_g] = z\} &= \\ &= \sum_{\chi \in \text{Irr}(G)} \frac{|G|^{2g-2} \chi(z)}{\chi(1)^{2g-2} \chi(1)} \end{aligned}$$



Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

Let  $z \in G$  in a finite group  $G$  then

$$\begin{aligned} \frac{1}{|G|} \#\{a_j, b_j \in G \mid [a_1, b_1] \dots [a_g, b_g] = z\} &= \\ &= \sum_{\chi \in \text{Irr}(G)} \frac{|G|^{2g-2} \chi(z)}{\chi(1)^{2g-2} \chi(1)} \end{aligned}$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q)$$

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

Let  $z \in G$  in a finite group  $G$  then

$$\begin{aligned} \frac{1}{|G|} \#\{a_j, b_j \in G \mid [a_1, b_1] \dots [a_g, b_g] = z\} &= \\ &= \sum_{\chi \in \text{Irr}(G)} \frac{|G|^{2g-2} \chi(z)}{\chi(1)^{2g-2} \chi(1)} \end{aligned}$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) \stackrel{\text{Katz}}{=} |\mathcal{M}_B^n(\mathbb{F}_q)|$$

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

Let  $z \in G$  in a finite group  $G$  then

$$\begin{aligned} \frac{1}{|G|} \#\{a_j, b_j \in G \mid [a_1, b_1] \dots [a_g, b_g] = z\} &= \\ &= \sum_{\chi \in \text{Irr}(G)} \frac{|G|^{2g-2} \chi(z)}{\chi(1)^{2g-2} \chi(1)} \end{aligned}$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) \stackrel{\text{Katz}}{=} |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2} (q-1) \chi(\xi_n^d)}{\chi(1)^{2g-2} \chi(1)}$$

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

Let  $z \in G$  in a finite group  $G$  then

$$\begin{aligned} \frac{1}{|G|} \#\{a_j, b_j \in G \mid [a_1, b_1] \dots [a_g, b_g] = z\} &= \\ &= \sum_{\chi \in \text{Irr}(G)} \frac{|G|^{2g-2} \chi(z)}{\chi(1)^{2g-2} \chi(1)} \end{aligned}$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) \stackrel{\text{Katz}}{=} |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2} (q-1) \chi(\xi_n^d)}{\chi(1)^{2g-2} \chi(1)}$$

- e.g.  $\frac{E(\mathcal{M}_B^2)}{(q-1)^{2g}}$

Theorem (Frobenius 1896, Hurwitz 1902, Serre 1992, Freed-Quinn 1993,...)

Let  $z \in G$  in a finite group  $G$  then

$$\begin{aligned} \frac{1}{|G|} \#\{a_j, b_j \in G \mid [a_1, b_1] \dots [a_g, b_g] = z\} &= \\ &= \sum_{\chi \in \text{Irr}(G)} \frac{|G|^{2g-2} \chi(z)}{\chi(1)^{2g-2} \chi(1)} \end{aligned}$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) \stackrel{\text{Katz}}{=} |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2} (q-1) \chi(\xi_n^d)}{\chi(1)^{2g-2} \chi(1)}$$

- e.g.  $\frac{E(\mathcal{M}_B^2)}{(q-1)^{2g}} =$

$$(q^2-1)^{2g-2} + q^{2g-2}(q^2-1)^{2g-2} - \frac{1}{2}q^{2g-2}(q-1)^{2g-2} - \frac{1}{2}q^{2g-2}(q+1)^{2g-2}$$

# Weight polynomials of $\mathcal{M}_B^n$

Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

# Weight polynomials of $\mathcal{M}_B^n$

Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

# Weight polynomials of $\mathcal{M}_B^n$

Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\frac{WH(\mathcal{M}_B^2; q, t)}{(1 + qt)^{2g}}$$



Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\begin{aligned} \frac{WH(\mathcal{M}_B^2; q, t)}{(1 + qt)^{2g}} &= \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \\ &- \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)} \end{aligned}$$

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\begin{aligned} \frac{WH(\mathcal{M}_B^2; q, t)}{(1 + qt)^{2g}} &= \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \\ &- \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)} \end{aligned}$$

- Conjecture is consistent with formula for

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\begin{aligned} \frac{WH(\mathcal{M}_B^2; q, t)}{(1 + qt)^{2g}} &= \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \\ &- \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)} \end{aligned}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\begin{aligned} \frac{WH(\mathcal{M}_B^2; q, t)}{(1 + qt)^{2g}} &= \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \\ &- \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)} \end{aligned}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)
  - $P(\mathcal{M}_{\text{Dol}}^4; t)$  of (Garcia-Prada-Heinloth-Schmitt 2011)

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k} - 1)(1 - w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\frac{WH(\mathcal{M}_B^2; q, t)}{(1 + qt)^{2g}} = \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)
  - $P(\mathcal{M}_{\text{Dol}}^4; t)$  of (Garcia-Prada-Heinloth-Schmitt 2011)
  - $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Schiffmann, 2016) and (Mellit, 2017)

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k}-1)(1-w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\frac{WH(\mathcal{M}_B^2; q, t)}{(1+qt)^{2g}} = \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)
  - $P(\mathcal{M}_{\text{Dol}}^4; t)$  of (Garcia-Prada-Heinloth-Schmitt 2011)
  - $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Schiffmann, 2016) and (Mellit, 2017)
  - conjectured  $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Chuang-Diaconescu-Pan 2010)

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k}-1)(1-w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\frac{WH(\mathcal{M}_B^2; q, t)}{(1+qt)^{2g}} = \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)
  - $P(\mathcal{M}_{\text{Dol}}^4; t)$  of (Garcia-Prada-Heinloth-Schmitt 2011)
  - $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Schiffmann, 2016) and (Mellit, 2017)
  - conjectured  $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Chuang-Diaconescu-Pan 2010) via (Mozgovoy 2011)

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k}-1)(1-w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\frac{WH(\mathcal{M}_B^2; q, t)}{(1+qt)^{2g}} = \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)
  - $P(\mathcal{M}_{\text{Dol}}^4; t)$  of (Garcia-Prada-Heinloth-Schmitt 2011)
  - $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Schiffmann, 2016) and (Mellit, 2017)
  - conjectured  $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Chuang-Diaconescu-Pan 2010) via (Mozgovoy 2011) and (Mellit 2016)



## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k}-1)(1-w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\frac{WH(\mathcal{M}_B^2; q, t)}{(1+qt)^{2g}} = \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)
  - $P(\mathcal{M}_{\text{Dol}}^4; t)$  of (Garcia-Prada-Heinloth-Schmitt 2011)
  - $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Schiffmann, 2016) and (Mellit, 2017)
  - conjectured  $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Chuang-Diaconescu-Pan 2010) via (Mozgovoy 2011) and (Mellit 2016)
  - $WH(\mathcal{M}_{\text{Dol}}^n; q, t)$  by (Chuang-Diaconescu-Pan 2012) as refined Gopakumar-Vafa conj for local curve CY 3-fold

## Conjecture (Hausel-Villegas, 2008)

$$\sum_{\lambda} \prod \frac{(z^{2l+1} - w^{2a+1})^{2g}}{(z^{2l+2} - w^{2a})(z^{2l} - w^{2a+2})} T^{|\lambda|} = \exp \left( \sum_{n,k} \frac{WH(\mathcal{M}_B^n; w^{2k}, -(zw)^{-2k})(zw)^{dn}}{(z^{2k}-1)(1-w^{2k})} \frac{T^{nk}}{k} \right)$$

- when  $r = 2$  Conjecture  $\leadsto$  Theorem (Hausel-Villegas, 2008)

$$\frac{WH(\mathcal{M}_B^2; q, t)}{(1 + qt)^{2g}} = \frac{(q^2 t^3 + 1)^{2g}}{(q^2 t^2 - 1)(q^2 t^4 - 1)} + \frac{q^{2g-2} t^{4g-4} (q^2 t + 1)^{2g}}{(q^2 - 1)(q^2 t^2 - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt + 1)^{2g}}{(qt^2 - 1)(q - 1)} - \frac{1}{2} \frac{q^{2g-2} t^{4g-4} (qt - 1)^{2g}}{(q + 1)(qt^2 + 1)}$$

- Conjecture is consistent with formula for
  - $P(\mathcal{M}_{\text{Dol}}^3; t)$  of (Gothen 1994)
  - $P(\mathcal{M}_{\text{Dol}}^4; t)$  of (Garcia-Prada-Heinloth-Schmitt 2011)
  - $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Schiffmann, 2016) and (Mellit, 2017)
  - conjectured  $P(\mathcal{M}_{\text{Dol}}^n; t)$  by (Chuang-Diaconescu-Pan 2010) via (Mozgovoy 2011) and (Mellit 2016)
  - $WH(\mathcal{M}_{\text{Dol}}^n; q, t)$  by (Chuang-Diaconescu-Pan 2012) as refined Gopakumar-Vafa conj for local curve CY 3-fold via "P=W"



- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| =$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_B^n; q) = q^{d_n} E(\mathcal{M}_B^n; 1/q)$  palindromic

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_{\mathbb{B}}^n; q) = |\mathcal{M}_{\mathbb{B}}^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_{\mathbb{B}}^n; q) = q^{d_n} E(\mathcal{M}_{\mathbb{B}}^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_B^n; q) = q^{d_n} E(\mathcal{M}_B^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$



# Curious Hard Lefschetz

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_B^n; q) = q^{d_n} E(\mathcal{M}_B^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$

- $\leadsto$  Curious Hard Lefschetz Conjecture

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_B^n; q) = q^{d_n} E(\mathcal{M}_B^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$

- $\leadsto$  Curious Hard Lefschetz Conjecture (theorem when  $n = 2$ ):

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_B^n; q) = q^{d_n} E(\mathcal{M}_B^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$

- $\rightsquigarrow$  Curious Hard Lefschetz Conjecture (theorem when  $n = 2$ ):

$$L^l : Gr_{d_n-2l}^W H^{i-l}(\mathcal{M}_B^n) \xrightarrow{\cong} Gr_{d_n+2l}^W H^{i+l}(\mathcal{M}_B^n)$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_B^n; q) = q^{d_n} E(\mathcal{M}_B^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$

- $\rightsquigarrow$  Curious Hard Lefschetz Conjecture (theorem when  $n = 2$ ):

$$L^l : \begin{array}{ccc} \text{Gr}_{d_n-2l}^W H^{i-l}(\mathcal{M}_B^n) & \xrightarrow{\cong} & \text{Gr}_{d_n+2l}^W H^{i+l}(\mathcal{M}_B^n) \\ X & \mapsto & X \cup \alpha^l \end{array}$$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_{\mathbb{B}}^n; q) = |\mathcal{M}_{\mathbb{B}}^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_{\mathbb{B}}^n; q) = q^{d_n} E(\mathcal{M}_{\mathbb{B}}^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$

- $\leadsto$  Curious Hard Lefschetz Conjecture (theorem when  $n = 2$ ):

$$L^l : \begin{array}{ccc} \text{Gr}_{d_n-2l}^W H^{i-l}(\mathcal{M}_{\mathbb{B}}^n) & \xrightarrow{\cong} & \text{Gr}_{d_n+2l}^W H^{i+l}(\mathcal{M}_{\mathbb{B}}^n) \\ X & \mapsto & X \cup \alpha^l \end{array}$$

where  $\alpha \in W_4 H^2(\mathcal{M}_{\mathbb{B}}^n)$

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_B^n; q) = |\mathcal{M}_B^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_B^n; q) = q^{d_n} E(\mathcal{M}_B^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$

- $\rightsquigarrow$  Curious Hard Lefschetz Conjecture (theorem when  $n = 2$ ):

$$L^l : \underset{X}{\text{Gr}_{d_n-2l}^W H^{i-l}(\mathcal{M}_B^n)} \xrightarrow{\cong} \underset{X \cup \alpha^l}{\text{Gr}_{d_n+2l}^W H^{i+l}(\mathcal{M}_B^n)}$$

where  $\alpha \in W_4 H^2(\mathcal{M}_B^n)$

- The implied functional equation on the conjectured  
 $WH(\mathcal{M}_B^n; q, t) = (qt)^{d_n} WH(\mathcal{M}_B^n; \frac{1}{qt^2}, t)$  holds

- (Hausel-Villegas 2008) calculates

$$E(\mathcal{M}_{\mathbb{B}}^n; q) = |\mathcal{M}_{\mathbb{B}}^n(\mathbb{F}_q)| = \sum_{\chi \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))} \frac{|\text{GL}_n(\mathbb{F}_q)|^{2g-2}}{\chi(1)^{2g-2}} \frac{\chi(\xi_n^d)}{\chi(1)}$$

- we find  $E(\mathcal{M}_{\mathbb{B}}^n; q) = q^{d_n} E(\mathcal{M}_{\mathbb{B}}^n; 1/q)$  palindromic  
by *Alvis-Curtis duality*

$$q^{\frac{n(n-1)}{2}} \chi(1)(1/q) = \chi'(1)(q) \text{ for dual pair } \chi, \chi' \in \text{Irr}(\text{GL}_n(\mathbb{F}_q))$$

- $\rightsquigarrow$  Curious Hard Lefschetz Conjecture (theorem when  $n = 2$ ):

$$L^l : \underset{X}{\text{Gr}_{d_n-2l}^W H^{i-l}(\mathcal{M}_{\mathbb{B}}^n)} \xrightarrow{\cong} \underset{X \cup \alpha^l}{\text{Gr}_{d_n+2l}^W H^{i+l}(\mathcal{M}_{\mathbb{B}}^n)}$$

where  $\alpha \in W_4 H^2(\mathcal{M}_{\mathbb{B}}^n)$

- The implied functional equation on the conjectured  $WH(\mathcal{M}_{\mathbb{B}}^n; q, t) = (qt)^{d_n} WH(\mathcal{M}_{\mathbb{B}}^n; \frac{1}{qt^2}, t)$  holds
- What is the meaning of  $W$  on  $H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\mathbb{B}}^n)$ ?

# Perverse filtration



# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  
 $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves
- recipe (de Cataldo-Migliorini, 2008) for perverse filtration when  $X$  smooth and  $Y$  affine:

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves
- recipe (de Cataldo-Migliorini, 2008) for perverse filtration when  $X$  smooth and  $Y$  affine:  
$$Y_0 \subset \dots \subset Y_i \subset \dots \subset Y_n = Y$$

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves
- recipe (de Cataldo-Migliorini, 2008) for perverse filtration when  $X$  smooth and  $Y$  affine:  
 $Y_0 \subset \dots \subset Y_i \subset \dots \subset Y_n = Y$  s.t.  $Y_i$  generic with  $\dim(Y_i) = i$



# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves
- recipe (de Cataldo-Migliorini, 2008) for perverse filtration when  $X$  smooth and  $Y$  affine:  
 $Y_0 \subset \dots \subset Y_i \subset \dots \subset Y_n = Y$  s.t.  $Y_i$  generic with  $\dim(Y_i) = i$

$$P_{k-i-1}H^k(X) = \ker(H^k(X) \rightarrow H^k(f^{-1}(Y_i)))$$

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves
- recipe (de Cataldo-Migliorini, 2008) for perverse filtration when  $X$  smooth and  $Y$  affine:  
 $Y_0 \subset \dots \subset Y_i \subset \dots \subset Y_n = Y$  s.t.  $Y_i$  generic with  $\dim(Y_i) = i$

$$P_{k-i-1}H^k(X) = \ker(H^k(X) \rightarrow H^k(f^{-1}(Y_i)))$$

- the Relative Hard Lefschetz Theorem holds:

$$L^l : Gr_{d-l}^P H^*(X) \xrightarrow{\cong} Gr_{d+l}^P H^{*+2l}(X)$$

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves
- recipe (de Cataldo-Migliorini, 2008) for perverse filtration when  $X$  smooth and  $Y$  affine:  
 $Y_0 \subset \dots \subset Y_i \subset \dots \subset Y_n = Y$  s.t.  $Y_i$  generic with  $\dim(Y_i) = i$

$$P_{k-i-1}H^k(X) = \ker(H^k(X) \rightarrow H^k(f^{-1}(Y_i)))$$

- the Relative Hard Lefschetz Theorem holds:

$$L^l : \begin{array}{ccc} Gr_{d-l}^P H^*(X) & \xrightarrow{\cong} & Gr_{d+l}^P H^{*+2l}(X) \\ X & \mapsto & X \cup \alpha^l \end{array}$$

# Perverse filtration

- $f : X \rightarrow Y$  a *proper* map between complex algebraic varieties of relative dimension  $d$
- (de Cataldo-Migliorini 2005) introduce *perverse filtration*  $P_0 \subset \dots \subset P_i \subset \dots \subset P_k(X) \cong H^k(X)$  from the study of the Beilinson-Bernstein-Deligne-Gabber decomposition theorem for  $Rf_*(\mathbb{Q}_X)$  into perverse sheaves
- recipe (de Cataldo-Migliorini, 2008) for perverse filtration when  $X$  smooth and  $Y$  affine:  
 $Y_0 \subset \dots \subset Y_i \subset \dots \subset Y_n = Y$  s.t.  $Y_i$  generic with  $\dim(Y_i) = i$

$$P_{k-i-1}H^k(X) = \ker(H^k(X) \rightarrow H^k(f^{-1}(Y_i)))$$

- the Relative Hard Lefschetz Theorem holds:

$$L^l : \begin{array}{ccc} Gr_{d-l}^P H^*(X) & \xrightarrow{\cong} & Gr_{d+l}^P H^{*+2l}(X) \\ X & \mapsto & X \cup \alpha^l \end{array}$$

where  $\alpha \in H^2(X)$  is a relative ample class

# P=W conjecture

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*



# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$$

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n) \text{ under the isomorphism}$$
$$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$$

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

Theorem (de Cataldo-Hausel-Migliorini 2010)

$P = W$  for  $n = 2$

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

Theorem (de Cataldo-Hausel-Migliorini 2010)

$P = W$  for  $n = 2$

- (de Cataldo-Hausel-Migliorini 2011)  $\rightsquigarrow P = W$

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

Theorem (de Cataldo-Hausel-Migliorini 2010)

$P = W$  for  $n = 2$

- (de Cataldo-Hausel-Migliorini 2011)  $\leadsto P = W$  when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are isomorphic to  $T^*E^{[n]}$  and  $(\mathbb{C}^\times \times \mathbb{C}^\times)^{[n]}$

# P=W conjecture

- recall Hitchin map  $\chi : \begin{array}{l} \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i) \\ (E, \phi) \mapsto \text{charpol}(\phi) \end{array}$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

Theorem (de Cataldo-Hausel-Migliorini 2010)

$P = W$  for  $n = 2$

- (de Cataldo-Hausel-Migliorini 2011)  $\sim P = W$  when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are isomorphic to  $T^*E^{[n]}$  and  $(\mathbb{C}^\times \times \mathbb{C}^\times)^{[n]}$
- (Shen, Zhang 2018)  $P = W$  holds in more parabolic cases of (Gröchenig 2014) when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are Hilbert schemes

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

Theorem (de Cataldo-Hausel-Migliorini 2010)

$P = W$  for  $n = 2$

- (de Cataldo-Hausel-Migliorini 2011)  $\sim P = W$  when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are isomorphic to  $T^*E^{[n]}$  and  $(\mathbb{C}^\times \times \mathbb{C}^\times)^{[n]}$
- (Shen, Zhang 2018)  $P = W$  holds in more parabolic cases of (Gröchenig 2014) when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are Hilbert schemes
- (Szabó, 2018) wild  $P = W$  holds in the Painlevé cases



# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

Theorem (de Cataldo-Hausel-Migliorini 2010)

$P = W$  for  $n = 2$

- (de Cataldo-Hausel-Migliorini 2011)  $\sim P = W$  when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are isomorphic to  $T^*E^{[n]}$  and  $(\mathbb{C}^\times \times \mathbb{C}^\times)^{[n]}$
- (Shen, Zhang 2018)  $P = W$  holds in more parabolic cases of (Gröchenig 2014) when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are Hilbert schemes
- (Szabó, 2018) wild  $P = W$  holds in the Painlevé cases
- (de Cataldo, Maulik 2018)  $P$  is topological

# P=W conjecture

- recall Hitchin map  $\chi : \mathcal{M}_{\text{Dol}}^n \rightarrow \mathbb{A} := \bigoplus_{i=1}^n H^0(C; K^i)$   
 $(E, \phi) \mapsto \text{charpol}(\phi)$
- (Hitchin 1987)  $\rightarrow$  completely integrable Hamiltonian system and *proper*

Conjecture ("P=W", de Cataldo-Hausel-Migliorini 2008)

$P_k(\mathcal{M}_{\text{Dol}}^n) \cong W_{2k}(\mathcal{M}_{\text{B}}^n)$  under the isomorphism

$H^*(\mathcal{M}_{\text{Dol}}^n) \cong H^*(\mathcal{M}_{\text{B}}^n)$  from non-Abelian Hodge theory

Theorem (de Cataldo-Hausel-Migliorini 2010)

$P = W$  for  $n = 2$

- (de Cataldo-Hausel-Migliorini 2011)  $\sim P = W$  when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are isomorphic to  $T^*E^{[n]}$  and  $(\mathbb{C}^\times \times \mathbb{C}^\times)^{[n]}$
- (Shen, Zhang 2018)  $P = W$  holds in more parabolic cases of (Gröchenig 2014) when  $\mathcal{M}_{\text{Dol}}$  and  $\mathcal{M}_{\text{B}}$  are Hilbert schemes
- (Szabó, 2018) wild  $P = W$  holds in the Painlevé cases
- (de Cataldo, Maulik 2018)  $P$  is topological  $\Leftarrow P = W$

# Mirror symmetry and $P = W$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*

# Mirror symmetry and $P = W$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$

# Mirror symmetry and $P = W$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$
- (Hausel 2010) conjectures  $H^{i,j}(\text{Gr}_p^P H^{i+j}(\mathcal{M}_{\text{Dol}}(\text{SL}_n))) \cong H^{j+d/2-p, i+d/2-p}(\text{Gr}_{d-p}^P H_{\text{st}}^{i+j+d-2p}(\mathcal{M}_{\text{Dol}}(\text{PGL}_n)))$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\leadsto H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$
- (Hausel 2010) conjectures  $H^{i,j}(\text{Gr}_p^P H^{i+j}(\mathcal{M}_{\text{Dol}}(\text{SL}_n))) \cong H^{j+d/2-p, i+d/2-p}(\text{Gr}_{d-p}^P H_{\text{st}}^{i+j+d-2p}(\mathcal{M}_{\text{Dol}}(\text{PGL}_n)))$
- (Gröchenig, Wyss, Ziegler 2017) proves topological mirror symmetry using  $p$ -adic integration along the Hitchin map



- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$
- (Hausel 2010) conjectures  $H^{i,j}(\text{Gr}_p^P H^{i+j}(\mathcal{M}_{\text{Dol}}(\text{SL}_n))) \cong H^{j+d/2-p, i+d/2-p}(\text{Gr}_{d-p}^P H_{\text{st}}^{i+j+d-2p}(\mathcal{M}_{\text{Dol}}(\text{PGL}_n)))$
- (Gröchenig, Wyss, Ziegler 2017) proves topological mirror symmetry using  $p$ -adic integration along the Hitchin map
- (Gröchenig, Wyss, Ziegler 2018)  $\rightsquigarrow$  Ngô's geometric stabilization, implying the Fundamental Lemma

# Mirror symmetry and $P = W$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$
- (Hausel 2010) conjectures  $H^{i,j}(\text{Gr}_p^P H^{i+j}(\mathcal{M}_{\text{Dol}}(\text{SL}_n))) \cong H^{i+d/2-p,j+d/2-p}(\text{Gr}_{d-p}^P H_{\text{st}}^{i+j+d-2p}(\mathcal{M}_{\text{Dol}}(\text{PGL}_n)))$
- (Gröchenig, Wyss, Ziegler 2017) proves topological mirror symmetry using  $p$ -adic integration along the Hitchin map
- (Gröchenig, Wyss, Ziegler 2018)  $\rightsquigarrow$  Ngô's geometric stabilization, implying the Fundamental Lemma
- $P = W$  and topological mirror symmetry  $\rightsquigarrow$

# Mirror symmetry and $P = W$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$
- (Hausel 2010) conjectures  $H^{i,j}(\text{Gr}_p^P H^{i+j}(\mathcal{M}_{\text{Dol}}(\text{SL}_n))) \cong H^{i+d/2-p,j+d/2-p}(\text{Gr}_{d-p}^P H_{\text{st}}^{i+j+d-2p}(\mathcal{M}_{\text{Dol}}(\text{PGL}_n)))$
- (Gröchenig, Wyss, Ziegler 2017) proves topological mirror symmetry using  $p$ -adic integration along the Hitchin map
- (Gröchenig, Wyss, Ziegler 2018)  $\rightsquigarrow$  Ngô's geometric stabilization, implying the Fundamental Lemma
- $P = W$  and topological mirror symmetry  $\rightsquigarrow E(\mathcal{M}_{\text{B}}(\text{SL}_n)); q \cong E_{\text{st}}(\mathcal{M}_{\text{B}}(\text{PGL}_n))$  (Hausel, Mereb, Villegas 2019)

# Mirror symmetry and $P = W$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$
- (Hausel 2010) conjectures  $H^{i,j}(\text{Gr}_p^P H^{i+j}(\mathcal{M}_{\text{Dol}}(\text{SL}_n))) \cong H^{i+d/2-p,j+d/2-p}(\text{Gr}_{d-p}^P H_{\text{st}}^{i+j+d-2p}(\mathcal{M}_{\text{Dol}}(\text{PGL}_n)))$
- (Gröchenig, Wyss, Ziegler 2017) proves topological mirror symmetry using  $p$ -adic integration along the Hitchin map
- (Gröchenig, Wyss, Ziegler 2018)  $\rightsquigarrow$  Ngô's geometric stabilization, implying the Fundamental Lemma
- $P = W$  and topological mirror symmetry  $\rightsquigarrow E(\mathcal{M}_{\text{B}}(\text{SL}_n)); q \cong E_{\text{st}}(\mathcal{M}_{\text{B}}(\text{PGL}_n))$  (Hausel, Mereb, Villegas 2019) from character table of  $\text{SL}_n(\mathbb{F}_q)$

# Mirror symmetry and $P = W$

- $Y$  and  $\tilde{Y}$  mirror symmetric CY 3-folds  $\rightsquigarrow H^{i,j}(Y) \cong H^{d_Y-i,j}(\tilde{Y})$   
*topological mirror test*
- (Hausel-Thaddeus 2002) conjectures topological mirror symmetry  $H^k(\mathcal{M}_{\text{Dol}}(\text{SL}_n)) \cong H_{\text{st}}^k(\mathcal{M}_{\text{Dol}}(\text{PGL}_n))$
- (Hausel 2010) conjectures  $H^{i,j}(\text{Gr}_p^P H^{i+j}(\mathcal{M}_{\text{Dol}}(\text{SL}_n))) \cong H^{j+d/2-p, i+d/2-p}(\text{Gr}_{d-p}^P H_{\text{st}}^{i+j+d-2p}(\mathcal{M}_{\text{Dol}}(\text{PGL}_n)))$
- (Gröchenig, Wyss, Ziegler 2017) proves topological mirror symmetry using  $p$ -adic integration along the Hitchin map
- (Gröchenig, Wyss, Ziegler 2018)  $\rightsquigarrow$  Ngô's geometric stabilization, implying the Fundamental Lemma
- $P = W$  and topological mirror symmetry  $\rightsquigarrow E(\mathcal{M}_{\text{B}}(\text{SL}_n)); q \cong E_{\text{st}}(\mathcal{M}_{\text{B}}(\text{PGL}_n))$  (Hausel, Mereb, Villegas 2019) from character table of  $\text{SL}_n(\mathbb{F}_q)$
- (Katzarkov, Kontsevich, Pantev 2017) conjecture a version of  $P = W$  topological mirror symmetry for Landau-Ginzburg models