

10.2

Find an explicit formula for a_n :

5. $2, \frac{5}{2}, \frac{10}{3}, \frac{17}{4}, \frac{26}{5}, \dots$

6. $-\frac{1}{4}, \frac{2}{9}, -\frac{3}{16}, \frac{4}{25}, -\frac{5}{36}, \dots$

Determine the boundedness and monotonicity of each sequence:

29. $\{\frac{2^n-1}{2^n}\}$

30. $\{\frac{1}{2n} - \frac{1}{2n+3}\}$

A sequence $\{a_n\}$ is said to be defined recursively if, for some $k \geq 1$, the terms a_1, a_2, \dots, a_k are given and a_n is specified in terms of a_1, a_2, \dots, a_{n-1} for each $n \geq k$. The formula specifying a_n for $n \geq k$ in terms of some (or all) of its predecessors is called a recurrence relation. Write down the first six terms of each sequence and then give the general formula for a_n .

51. $a_1 = 1; a_{n+1} = a_n + 2n + 1$

52. $a_1 = 1; a_{n+1} = 2a_n + 1$

61. Let r be a real number, $r \neq 0$. Define a sequence $\{S_n\}$ by:

$$S_1 = 1$$

$$S_2 = 1 + r$$

$$S_3 = 1 + r + r^2$$

.

.

.

$$S_n = 1 + r + r^2 + \dots + r^{n-1}$$

.

.

.

(a) Suppose $r = 1$. What is S_n for $n = 1, 2, 3, \dots$?

(b) Suppose $r \neq 1$. Find a formula for S_n that does not involve adding up the powers of r . HINT: Calculate $S_n - rS_n$.

62. Let $a_n = \frac{1}{n(n+1)}, n = 1, 2, 3, \dots$, and let $\{S_n\}$ be the sequence defined by

$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

.

.

.

$$S_n = a_1 + a_2 + a_3 + \dots + a_n$$

.

.

.

Find a formula for $S_n, n = 1, 2, 3, \dots$, that does not involve adding up the terms a_1, a_2, a_3, \dots . HINT: Use partial fractions to write $1/[k(k+1)]$ as the sum of two fractions.

10.3

State whether the sequence converges and, if it does, find the limit.

5. $\{\frac{n-1}{n}\}$

6. $\{\frac{n+(-1)^n}{n}\}$

31. $\{(1 + \frac{1}{n})^{2n}\}$

32. $\{(1 + \frac{1}{n})^{n/2}\}$

These sequences are defined recursively. Determine in each case whether the sequence converges and, if so, find the limit. Start each sequence with

$$a_1 = 1.$$

53. $a_{n+1} = \frac{1}{n+1}a_n$

54. $a_{n+1} = \frac{n}{n+1}a_n$