

INFERENCE FOR SIMPLE OLS

Model Assumptions ("The" Simple Linear Regression Model Version IV):

(We consider x_1, \dots, x_n as fixed.)

1. $E(Y|x) = \eta_0 + \eta_1 x$ (linear mean function)
2. $\text{Var}(Y|x) = \sigma^2$ (Equivalently, $\text{Var}(e|x) = \sigma^2$) (constant variance)
3. y_1, \dots, y_n are independent observations. (independence)
4. (NEW) $Y|x$ is normal for each x (normality)

(1) + (2) + (4) can be summarized as:

$$Y|x \sim N(\eta_0 + \eta_1 x, \sigma^2)$$

Recall: $e|x = Y|x - E(Y|x)$

So: $e|x \sim N(0, \sigma^2)$

i.e., all errors have the same distribution -- so we just say e instead of $e|x$.

Since $\hat{\eta}_0$ and $\hat{\eta}_1$ are linear combinations of the $Y|x_i$'s, (3) + (4) imply that $\hat{\eta}_0$ and $\hat{\eta}_1$ are normally distributed random variables (that is, their sampling distributions are normal).

Recalling that

$$E(\hat{\eta}_1) = \eta_1 \quad \text{Var}(\hat{\eta}_1) = \frac{\sigma^2}{SXX} \quad E(\hat{\eta}_0) = \eta_0 \quad \text{Var}(\hat{\eta}_0) = \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{SXX} \right),$$

We have

$$\hat{\eta}_1 \sim \hat{\eta}_0 \sim$$

Look more at $\hat{\eta}_1$: We can standardize to get

$$\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\sigma^2/SXX}} \sim N(0,1)$$

But we don't know σ^2 , so need to approximate it by $\hat{\sigma}^2$ -- in other words approximate

$\text{Var}(\hat{\eta}_1)$ by $\hat{\text{Var}}(\hat{\eta}_1) = [\text{s.e.}(\hat{\eta}_1)]^2 = \frac{\hat{\sigma}^2}{SXX}$. Thus we want to use $\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2/SXX}}$. But we can't

expect this to be normal, too. However,

$$\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2 / SXX}} =$$

$$(*) \quad \left(\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\sigma^2 / SXX}} \right) / \sqrt{\hat{\sigma}^2 / \sigma^2}$$

The numerator of (*) is normal (in fact, standard normal), as noted above.

Facts: (Proofs omitted)

a. $(n-2) \frac{\hat{\sigma}^2}{\sigma^2}$ has a χ^2 distribution with $n-2$ degrees of freedom

$$\text{Notation: } (n-2) \frac{\hat{\sigma}^2}{\sigma^2} \sim \chi^2(n-2)$$

b. $(n-2) \frac{\hat{\sigma}^2}{\sigma^2}$ is independent of $\hat{\eta}_1 - \eta_1$ (hence independent of the numerator in (*))

Comments on distributions:

1. A $\chi^2(k)$ distribution is defined as the distribution of a random variable which is a sum of squares of k independent standard normal random variables.

[Comment: Recall that $\hat{\sigma}^2 = \frac{1}{n-2} \text{RSS}$, so $(n-2) \frac{\hat{\sigma}^2}{\sigma^2} = \frac{\text{RSS}}{\sigma^2} = \sum \left(\frac{\hat{\epsilon}_i}{\sigma} \right)^2$ is a sum of n

squares; the fact quoted above says that it can also be expressed as a sum of $n-2$ squares of *independent standard normal* random variables.]

2. A t -distribution with k degrees of freedom is defined as the distribution of a random

variable of the form $\frac{Z}{\sqrt{U/k}}$ where

- $Z \sim N(0,1)$
- $U \sim \chi^2(k)$
- Z and U are independent.

In the fraction (*) above, take

$$U = (n-2) \frac{\hat{\sigma}^2}{\sigma^2} \sim \chi^2(n-2)$$

$$Z = \frac{\hat{\eta}_1 - \eta_1}{\sqrt{\sigma^2 / SXX}} \sim N(0,1)$$

Thus:
$$\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2 / SXX}} \sim t(n-2),$$

so we can do inference on η_1 , using $t = \frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2 / SXX}}$ as our test statistic.

Inference on η_0

With the same assumptions, it can be shown in an analogous manner (details omitted) that

$$\frac{\hat{\eta}_0 - \eta_0}{s.e.(\hat{\eta}_0)} \sim t(n-2),$$

so we can use this statistic to do inference on η_0 .