

Diophantine geometry in characteristic p : a survey

José Felipe Voloch

... it goes without saying that the function-fields over finite fields must be granted a fully simultaneous treatment with number-fields, instead of the segregated status, and at best the separate but equal facilities, which hitherto have been their lot. That, far from losing by such treatment, both races stand to gain by it,...

André Weil, 1967

1. Introduction

The purpose of this paper is to survey some of the recent finiteness results on rational and integral points on algebraic varieties defined over global fields of positive characteristics.

In the classical case of number fields, the basic results are the Mordell-Weil theorem, the Thue-Siegel-Roth theorem and Siegel's theorem on integral points on curves (see [L1]). Recently, there have been some major developments, first Faltings proved the Mordell conjecture, then Vojta gave another proof and this latter method was extended by Faltings [F3,4] to prove two outstanding conjectures of Lang. There are some historical comments in [F2] and [L2] is a general survey of the subject. In the number field case there are also very general quantitative conjectures made by Vojta [Vj]. These seem very deep and beyond the reach of current techniques. Of course, one should consider also the case of function fields of characteristic zero and there has been considerable progress there also. We refer to [L2] and [B2] for more details. In [B2] there is also a description of a tentative transposition of the function field methods to the number field case.

The situation in positive characteristic is similar to that of number fields. Results similar to those of Faltings have been proved recently, by different methods. We shall describe these results in detail below. One would hope to have conjectures similar to Vojta's in the function field case. However, the direct transposition of Vojta's conjectures to the case of positive characteristics is false. It also seems that any simple modification of the conjectures has counterexamples. We have been unable to formulate a plausible conjecture, even in the case of \mathbf{P}^1 . We will present below several examples that illustrate pathological behaviour in positive characteristic and perhaps a more perspicacious reader will find a general pattern in which these examples will fit.

We shall not attempt a full description of the history of the subject but let us point out a few highlights before proceeding to a description of recent results. The analogue of the Mordell-Weil theorem was proved by Lang and Néron (see [L1]) guided by Severi's remark that it was related to his Theorem of the Base. Diophantine approximation was considered by Mahler and Osgood initially but the results are still fragmentary. (See section 4). In particular, Mahler showed that Liouville's inequality cannot be improved in general and so Siegel's argument to deal with integral points on curves over number fields (which were successfully transferred to the case of function fields of characteristic zero by Lang) cannot be applied in characteristic p . It is unclear who first proved that elliptic curves in characteristic p with non-constant j -invariant have finitely many integral points. It seems to be well-known that this follows formally from the Mordell conjecture, (see e.g. [B1], ch 7, thm. 3.1) which was proved first by Samuel ([Sa]) but it is unclear who first noticed this formal consequence. The result was proved by Mason ([Ma], proof of thm. 14, pg. 114) and another proof given by the author [V4]. As mentioned above, the Mordell conjecture was proved by Samuel

([Sa1,2]), extending Grauert's proof for function fields of characteristic zero to characteristic p . Szpiro then gave an effective proof ([Sz]) as a consequence of his work on the Shafarevich conjecture. Let us also mention that recently Pheidas ([Ph]) proved that the problem of deciding if a polynomial in several variables and coefficients in $\mathbf{F}_q(t)$ has a zero with coordinates in $\mathbf{F}_q(t)$ is unsolvable. As usual, this has not abated the search for positive results in this area.

We do not, strictly speaking, follow Weil's advice in the above quotation. In fact, most of the recent success in the positive characteristic case is due to exploiting the special circumstances of this case.

A fundamental notion in this study is that of *isotriviality*. Roughly speaking, one is interested in deciding whether a variety defined over a function field can be defined over its constant field. We devote an appendix to this notion, studying it from different points of view.

Throughout this paper, K will denote a global field of positive characteristic p . In other words, K is a function field in one variable over a finite field of characteristic p .

We plan to update this survey as new results come to our attention. Updates will be posted in the author's home page in the World Wide Web. (URL: <http://www.ma.utexas.edu/users/voloch>)

2. Curves

For curves one would expect finiteness for rational/integral points provided the genus is large enough and the curve is non-isotrivial. In characteristic $p > 0$, already the notion of genus has a twist to it. Let C be an algebraic curve defined over a function field K of characteristic $p > 0$. One can define the (absolute) genus of C by extending the field to the algebraic closure. Another option is to define the genus of C relative to K to be

the integer g_K that makes the Riemann-Roch formula hold, that is, for any K -divisor D of C , of sufficiently large degree, the dimension, $l(D)$, of the K -vector space of functions of $K(C)$ whose polar divisor is bounded by D , is $\deg D + 1 - g_K$. Since K is not perfect, the relative genus may change under inseparable extensions. Luckily, this type of curve is actually easier to handle and it can be shown ([V3]) that if the genus of C relative to K is different from the (absolute) genus of C then $C(K)$ is finite. (This result was extended by S. T. Jeong, who obtained the same conclusion without the assumption that the constant field of K is finite.) We will now restrict our discussion to curves that do not change genus so there is now no loss of generality in assuming the curves to be smooth.

As mentioned above, Samuel proved the Mordell conjecture, to the effect that a curve of (absolute) genus at least two, which is non-isotrivial, has only finitely many rational points over a function field.

It remains to discuss integral points on *affine* curves. As expected,

Theorem 1. *Non-isotrivial affine curves (in the sense of the appendix) over function fields over finite fields have finitely many integral points.*

Sketch of proof: Let C be such a curve and \bar{C} is completion. Then there exists a cover X of \bar{C} branched over $\bar{C} \setminus C$ only, of degree prime to the characteristic, such that X is of genus at least 2. If \bar{C} has genus at least two, this is the Kodaira-Parshin construction ([L2],[Sz]), otherwise it is elementary. It also follows from the construction that X is non-isotrivial. Integral points on C over a fixed ring will then lift to rational points on X over a fixed field and the theorem then follows from the Mordell conjecture.

It is worth remarking at this point that the projective line minus three points is always isotrivial.

Another topic is to find bounds for the height of rational points on curves

of genus at least 2 (i.e. effective Mordell). Szpiro ([Sz]) had the first result on this line, which was improved by Moriwaki ([Mo]) and then by Kim ([Ki]) who obtained the following result:

Theorem 2 (Kim). *Let X be an algebraic curve of genus $g \geq 2$ defined over a function field K of characteristic $p > 0$, such that X is not defined over K^p . Then, for all points $P \in X$ with $K(P)/K$ finite one has, for $g > 2$,*

$$h_{K_X}(P) \leq (2g - 2)d(P) + O(h_{K_X}(P)^{1/2}),$$

and for $g = 2$,

$$h_{K_X}(P) \leq (2 + \epsilon)d(P) + O(1)$$

for any given $\epsilon > 0$. If moreover the Kodaira-Spencer class of X/K is of maximal rank (see appendix) then, for all $g \geq 2$:

$$h_{K_X}(P) \leq (2 + \epsilon)d(P) + O(1)$$

where $h_{K_X}(P)$ is the (logarithmic) height of P with respect to the canonical divisor K_X of X and $d(P) = (2g(K(P)) - 2)/[K(P) : K]$, where $g(K(P))$ denotes the genus of $K(P)$, that is, $d(P)$ is the logarithmic discriminant of P .

The bound $h_{K_X}(P) \leq (2g - 2)d(P) + O(h_{K_X}(P)^{1/2})$ is, in general, best possible. An example showing this can be constructed as follows. Let X_0 be a curve defined over the finite field \mathbf{F}_q with q elements, let F be the Frobenius map of X_0 , $K = \mathbf{F}_q(X_0)$ the function field of X_0 and $P \in X_0(K)$ the generic point. Let X be a cover of X_0 ramified only at P if the genus of X_0 is at least 2 and ramified at P and 0 if X_0 is an elliptic curve. To construct X one can use the Kodaira-Parshin construction. Now consider the points Q_m , say, of X which lift the points $F^m(P) \in X_0(K)$. It is not hard to show that $h(Q_m) - (2g - 2)d(Q_m)$ is proportional to $\#X_0(\mathbf{F}_{q^m}) - q^m + O(1)$, where g

is the genus of X . As the latter expression is $O(q^{m/2})$ and this estimate cannot be improved in general, we get that Kim's bound is best possible. This argument also shows that Kim's bound implies the Riemann hypothesis for curves over finite fields.

Another question is to bound the number of rational points. The following result was proved in [BV]:

Theorem 3. *Let C be an algebraic curve of genus at least 2 defined over a function field K of characteristic $p > 0$, such that C is not defined over K^p . Let J be the Jacobian of C , then*

$$\#C(K) \leq \#(J(K)/pJ(K)) \cdot p^g \cdot 3^g \cdot (8g - 2) \cdot g!.$$

The work of Caporaso et al. [CHM] shows that, in the number field case, uniform bounds for the number of rational points would follow from conjectures of Lang on varieties of general type (see below). In [AV2] similar consequences of these conjectures are obtained in the function field case. As it turns out, counterexamples can be given to some of these consequences in characteristic p ([AV2]). Namely, we show that there are no uniform bounds for the number of points on curves that change genus or to the number of separable maps between curves depending on the gonality of the curve.

We conclude this section by mentioning the work of Denis [D2], where he finds all rational points in a certain class of curves which, from the point of view of Drinfeld modules, are the characteristic p analogues of the Fermat curves. These curves change genus in the above sense.

3. Abelian varieties and their subvarieties

The following theorem was proved by Abramovich and the author [AV1], under restrictive hypotheses and then by Hrushovski [H], in general. It is the characteristic p analogue of a conjecture of Lang.

Theorem 3 (Hrushovski). *Let $X \subset A$ be a closed, integral subvariety of a semi abelian variety over an algebraically closed field of characteristic $p > 0$. Let $\Sigma \subset A$ be a subgroup of closed points such that $\text{rk}_{\mathbf{Z}_p} \Sigma \otimes \mathbf{Z}_p < \infty$. Let $\text{Stab}(X)$ be the stabilizer of X , that is the maximal subgroup-scheme of A leaving X invariant. If $\Sigma \cap X$ is Zariski dense in X then $X/\text{Stab}(X)$ is weakly isotrivial.*

Remark. The condition on the subgroup Σ holds if, e.g., it is contained in the prime-to- p division saturation of a finitely generated group. We conjecture that in this case we can include p -division. In this direction, Boxall [Bo] has shown that a similar result holds if Σ is a torsion group for which the orders of all elements are divisible only by a finite set of primes, in particular it holds for the p -power torsion subgroup. In a different vein, Denis [D1] has proposed some conjectures, analogous to the above, replacing abelian varieties by Drinfeld modules and their higher dimensional generalizations.

The case where X is a curve defined over a function field K , A its Jacobian and $\Sigma = A(K)$, the above theorem reduces to the Mordell conjecture which was proven by Grauert and Samuel (see [Sa1,2]).

Let A/K be an abelian variety of dimension n . For any closed subscheme $X \subset A$ there is a function $\lambda_v(X, \cdot) : A(K_v) \rightarrow [0, \infty]$ which satisfies the following property: for any affine open set $U \subset A$ and any system of generators $g_1, \dots, g_m \in \mathcal{O}(U)$ of the ideal defining $X \cap U$ in U , we may write $\lambda_v(X, P) = \min\{v(g_1(P)), \dots, v(g_m(P))\} + b(P)$ with b bounded on any bounded subset of $U(K_v)$. The function $\lambda_v(X, \cdot)$ is uniquely determined by the above property up to the addition of a bounded function and is called the local height function associated to X . This notion is developed in detail in [Si].

The characteristic p analogue of a conjecture of Lang predicts that if A is an abelian variety over a function field K of characteristic $p > 0$, the K/k -

trace of A is zero and X is an ample divisor on A then the set of integral points of $A \setminus X$ is finite. In trying to prove this conjecture, I was led to formulate an “infinitesimal” analogue of the Mordell-Lang conjecture. This was proved by Hrushovski. The following result is theorem 6.3 of [H].

Theorem 4 (Hrushovski). *Let A and X be as above and assume that the K/k -trace of A is zero. Then there exists a subvariety Y of X , defined over \bar{K} , which is a finite union of translates of abelian subvarieties of A , such that $\lambda_v(X, P) \ll \lambda_v(Y, P) + 1$ for all $P \in A(K)$.*

Using the above result and some estimates on distances between points on abelian varieties I managed to prove the following results in [V5]:

Theorem 5. *Let A be an ordinary abelian variety over a function field K of characteristic $p > 0$ and v a place of K and assume that the K/k -trace of A is zero and that $A[p^\infty] \cap A(K_s)$ is finite. Let X be a subvariety of A . Then $\lambda_v(X, P) \ll h(P)^{1/2}$ for all $P \in A(K), P \notin X$.*

Remark: The hypotheses of the theorem hold for sufficiently general A , see the appendix.

Corollary . *Hypotheses as in Theorem 5. Assume further that X is an ample divisor. Then for any finite set S of places of K , the set of S -integral points of $A \setminus X$ is finite.*

Proof of the corollary : In this case, $h(P)$, for an S -integral point of $A \setminus X$, is the sum of $\lambda_v(X, P)$ over the elements of S , and it follows that the height is bounded, which proves the corollary.

Example: Let A be a supersingular abelian variety, then $\lambda_v(\mathcal{O}, pP) = p^2 \lambda_v(\mathcal{O}, P)$ for P near \mathcal{O} , therefore, considering the sequence $p^n P$ for suitable P near \mathcal{O} , we get infinitely many points on $A(K)$ with $\lambda_v(\mathcal{O}, P) \gg h(P)$. It follows that for any subvariety X of A containing \mathcal{O} , we get $\lambda_v(X, P) \gg h(P)$.

Note that we can choose A to be nonisotrivial, although it is always going to be isogenous to isotrivial (see the appendix). However, we can choose X suitably so that $A \setminus X$ is not isotrivial.

4. Diophantine Approximation in characteristic p

In this section we will be concerned about the approximation of functions, algebraic over a global field K of positive characteristic by elements of K with respect to a valuation v of K . We define, for $y \in K_v \setminus K$ (although we will consider only y algebraic over K in what follows):

$$\alpha(y) = \limsup_{r \in K} v(y - r)/h(r),$$

where $h(r) = [K : k(r)]$, where k is the constant field of K . We will give some examples that exhibit pathological behaviour. Recall that $2 \leq \alpha(y) \leq d(y) := [K(y) : K]$, which are analogues of the classical theorems of Dirichlet and Liouville. Osgood [O] has shown that $\alpha(y) \leq [(d(y) + 1)/2]$ if y does not satisfy a Riccati equation and I proved the same bound if the cross ratio of any four conjugates of y over K is non constant. Finally, Lasjaunias and de Mathan [LdM] proved the same bound provided y does not satisfy $y^q = (ay + b)/(cy + d)$ where q is a power of the characteristic, $a, b, c, d \in K, ad - bc \neq 0$, which subsumes the previous results. There are some results on $\alpha(y)$ if y satisfies $y^q = (ay + b)/(cy + d)$ where $a, b, c, d \in K, ad - bc \neq 0$ and q is a power of p , due to the author [V1] and others ([BS],[dM],[MR]). We shall give examples that show that the bound of Lasjaunias and de Mathan is close to being best possible.

Take $K = k(x)$ and y satisfying $y^p - y = x$ and $z = y^2$ (y is a classical example of Mahler's). We have $\alpha(y) = d(y) = d(z) = p$. Also, whenever $v(y - r)/h(r)$ is near p we have $v(z - r^2)/h(r^2)$ near $p/2$. It follows that $\alpha(z) = p/2$. Note that z does not satisfy a Riccati equation. This example

can be generalized as follows: Given y and $R(Y) \in K(Y)$ a rational function of degree d in Y , then $d(R(y)) \leq d(y)$ and $\alpha(R(y)) \geq \alpha(y)/d$. So if $\alpha(y)$ is large we get new examples of well approximated functions which in general do not satisfy Riccati equations.

By definition, the cross ratio of x_1, \dots, x_4 is

$$[x_1, x_2, x_3, x_4] = (x_4 - x_1)(x_3 - x_2)/(x_4 - x_2)(x_3 - x_1).$$

Remark: Let D be the divisor on \mathbf{P}^1 formed by the conjugates of y over K , so D is of degree d and is defined over K . Let X be the affine curve $\mathbf{P}^1 \setminus D$. It can be checked that y satisfies a Riccati equation if and only if the Kodaira-Spencer class of X , in the sense of the appendix, vanishes. It can also be checked that the cross ratio of any four conjugates of y lies in k if and only if X is isotrivial, that is, isomorphic to an affine curve defined over k perhaps after field extension. It then follows from the results mentioned above that, when X is non-isotrivial, it has only finitely many integral points, which gives another proof of theorem 1 in the genus zero case.

Wang [W] has some results on diophantine approximation in \mathbf{P}^n for $n > 1$. In particular, she has a sufficient condition for finiteness of the set of integral points of \mathbf{P}^n minus $2n + 2$ hyperplanes.

5. Omitted topics

I felt I could not do justice to and give a proper survey on the following topics, which I'll just mention with a few references. I may have forgotten a few important ones and apologise to any reader whose favourite topic is not mentioned.

(a) Moduli of curves and abelian varieties. The work of Parshin, Zahrin and Szpiro led to the proof that there are only finitely many curves

of fixed genus over a fixed function field with a prescribed set of places of bad reduction and there are only finitely many abelian varieties over a fixed function field of characteristic p , prime-to- p isogenous to a given abelian variety. See [Sz] and [MB].

(b) Varieties of general type. Lang conjectured that, on a variety of general type over a number field, the set of rational points is not Zariski dense. One may try to transpose Lang's conjectures to the case of positive characteristic, but they are trivially false already in the case of curves. Two natural approaches to restore the conjecture, which work for curves, are either to insist on non - isotriviality of the variety or to look at points which are not in the image of the Frobenius map. Unfortunately both these approaches fail already for surfaces. In fact, there are unirational surfaces of general type in positive characteristic, and even non-constant families of those, which provide counterexamples to such conjectures. (See [AV2]). One may try to look at varieties with non-zero Kodaira - Spencer class, but there seem to be counterexamples here as well. Again the problem is due to unirational varieties. In all these examples the surfaces have a large set of birational endomorphisms (coming either from the Frobenius or from birational endomorphisms of \mathbf{P}^2), and one may try to take these into account in stating a Lang type conjecture. A rather drastic approach is to look only at varieties which are not covered by non-general type varieties, but this would be an unsatisfactory and almost unverifiable conjecture due to the fact that it not known how to tell whether a variety of general type can be covered by a variety which is not of general type.

In the positive direction, Martin-Deschamps and Lewin-Ménégaux [MDLM] proved that there are only finitely many separable dominating rational maps from X to Y , if X, Y are given varieties with Y of general type.

Another case of the conjecture is the following. Let A/K be an abelian variety and f a rational function on A . Consider the cover X of A defined by $z^p = f$. If A is defined over K^p then $X(K)$ is Zariski dense in X if and only if $f(P + P_0) \in K^p(A)$ for some $P_0 \in A(K)$, whereas if A is not defined over K^p this happens if and only if $f(V(P) + P_0) \in K^p(A^{(p)})$ for some $P_0 \in A(K)$, where $V : A^{(p)} \rightarrow A$ is the Verschiebung. A proof in the case of elliptic curves is given in [V3] and it readily generalizes to higher dimensions. The condition on f for $X(K)$ to be Zariski dense above is equivalent to vanishing of the Kodaira-Spencer class of X . Note that if A is simple of dimension at least 2 and f is not a p -th power, then X is of general type.

(c) The Birch and Swinnerton-Dyer conjecture. If A/K is an abelian variety, one defines an analytic function $L(E, s)$ and the conjecture of Birch and Swinnerton-Dyer states that the order of vanishing of $L(E, s)$ at $s = 1$ is the rank of $A(K)$ and gives a formula for the leading coefficient of the Taylor expansion of $L(E, s)$ around $s = 1$. Tate showed, for elliptic curves, that the first statement implies the second up to a power of p , which was removed by Milne, and that the conjecture was equivalent to the finiteness of the Tate-Shafarevich group. This has been generalized to higher dimensions, see [Mi].

(d) Existence of solutions to equations and inequalities We concentrated so far on finiteness statements, but one also expects that varieties which are "very rational" to have many rational points. For example, we have the Lang-Tsen theorem that if f_1, \dots, f_n are homogeneous polynomials over K of degrees d_1, \dots, d_n in at least $\sum d_i^2 + 1$ variables, then they have a common non-trivial zero. Carlitz generalized Tsen's method to deal with solutions of diophantine "inequalities" too. See [Ca], [Gre].

Appendix. Moduli, isotriviality, deformation theory and p -torsion points

Definition. Let X be a variety over a field K of characteristic $p > 0$. We say that X is *isotrivial* if after some base extension X is isomorphic to a variety X_1 that can be defined over a finite field. We say that X is *birationally isotrivial* if after some base extension X is birational to a variety X_1 that can be defined over a finite field. And we say that X is *weakly isotrivial* if after some base extension there is a rational map $X_1 \rightarrow X$, which induces a purely inseparable extension on the function fields, and such that X_1 can be defined over a finite field. To say that X_1 can be defined over a finite field means that there is Y/\mathbf{F}_q and a common extension L of K and \mathbf{F}_q such that $X_1 \otimes_K L$ is isomorphic to $Y \otimes_{\mathbf{F}_q} L$.

It is important to note that the notion of isotriviality is up to isomorphism. In particular it can happen that an affine open subset of an isotrivial variety is non-isotrivial. The notions of birationally isotrivial and weakly isotrivial, on the other hand, are birational notions. If K is the function field of a variety T defined over a finite field and X/K is a variety, then X corresponds to a family $\mathcal{X} \rightarrow T$ with generic fibre X . It is well-known that, in the projective case, X is isotrivial if and only if $\pi : \mathcal{X} \rightarrow T$ is generically constant, i.e., there is a non-empty open subset U of T such that the fibres over all points of U are isomorphic. (See e.g. [B1] Ch. 1, lemma (1.3)).

In characteristic zero, there is another equivalent notion of isotriviality, that of infinitesimally isotrivial. Although the equivalence of the notions does not hold in characteristic p , it is still very useful to consider infinitesimal isotriviality. To do that we first must define the Kodaira-Spencer class, in a more general setting than usual, following Katz [K].

Let \bar{X} be a smooth projective variety over K , D a divisor with normal

crossings on \bar{X} and $X = \bar{X} \setminus D$. Define a sheaf τ_X on \bar{X} of vector fields tangent to D , which is a subsheaf of the tangent sheaf of \bar{X} . Equivalently, τ_X is the sheaf of K -derivations of $\mathcal{O}_{\bar{X}}$ that preserve the ideal sheaf of D . If δ is a derivation on K we define the Kodaira-Spencer class $\kappa(\delta)$ in $H^1(\bar{X}, \tau_X)$ as follows. Let $\{U_i\}$ be an open cover of \bar{X} fine enough so that we can lift δ to derivations δ_i of $\mathcal{O}_{\bar{X}}(U_i)$, preserving the ideal of $D \cap U_i$. The 1-cocycle $\delta_i - \delta_j$ then defines the class $\kappa(\delta)$. We define X to be *infinitesimally isotrivial* if $\kappa(\delta) = 0$ for all δ .

Let X be a variety over a function field K of transcendence degree 1 over a finite field. It is convenient to study whether or not X is defined over $K^{(p)}$ using derivations: let t be a separating variable in K , and let C be an affine model of K over which $\delta = \partial/\partial t$ is a regular vector field. Let \mathcal{X} be a model of X , proper over C . Then that X can be defined over $K^{(p)}$ if and only if the derivation δ lifts to a vector field δ' over the inverse image in U of some open subset of C , satisfying $\delta'^p = 0$. See [V2], lemma 1 or [Og], lemma 3.5. Note that, obviously, δ lifts if and only if $\kappa(\delta) = 0$ and, as shown in loc. cit., the condition $\delta'^p = 0$ is automatically satisfied if $H^0(\bar{X}, \tau_X) = 0$.

Example: Let $A = E^g$ where E is a supersingular elliptic curve with $g \geq 2$. Choose a K -rational subspace of the Lie algebra of A not defined over K^p and corresponding to a height one group-scheme G . Take the quotient A/G , which is a non-isotrivial abelian variety over K with good reduction everywhere. If $K = k(t)$, then we can base change by any $k(s)/k(t)$, and obtain infinitely many families of such varieties. (See [Sz]).

For any projective smooth variety X , there is an obvious map

$$H^1(X, T_X) \rightarrow \text{Hom}(H^0(X, \Omega_X^1), H^1(X, \mathcal{O}_X)).$$

In the case of abelian varieties, this map is an isomorphism and we can talk about the rank of the Kodaira-Spencer class as a linear map between

vector spaces. In particular, we talk about the Kodaira-Spencer class having maximal rank, in this context. To justify the assertion, made in section 3, that abelian varieties with sufficiently general moduli satisfy the hypotheses of Theorem 5, we have the following result, proved in [V5].

Proposition. *Let A be an ordinary abelian variety over a function field K of characteristic $p > 0$ such that the Kodaira-Spencer map has maximal rank, then $A[p] \cap A(K_s) = \mathcal{O}$.*

Acknowledgements: The author would like to thank D. Abramovich for many helpful comments and the NSF (grant DMS-9301157) and the Alfred P. Sloan Foundation for financial support.

References.

- [AV1] D. Abramovich and J. F. Voloch, *Toward a proof of the Mordell- Lang conjecture in characteristic p* , International Math. Research Notices No. 5 (1992) 103-115.
- [AV2] D. Abramovich and J. F. Voloch, *Lang's conjectures, fibered powers, and uniformity*, New York Journal of Math., 2 (1996) 20-34.
- [BS] L. E. Baum and M. M. Sweet, *Badly approximable power series in characteristic 2*, Ann. Math. **105** (1977), 573-580.
- [Bo] J. Boxall, *Autour d'un probleme de Coleman* CRAS **315** (1992), 1063-1066.
- [B1] A. Buium, *Differential Algebra and Diophantine Geometry*, Hermann, Paris, 1994.
- [B2] A. Buium, *Differential Algebraic Geometry and Diophantine Geometry : an overview*, these proceedings.

- [BV] A. Buium, J. F. Voloch, *Mordell's conjecture in characteristic p : an explicit bound*, *Compositio Math.*, 103 (1996) 1-6.
- [Ca] L. Carlitz, *Some applications of a theorem of Chevalley*, *Duke Math. J.*, **18** (1951), 811 - 819.
- [CHM] L. Caporaso, J. Harris, B. Mazur, *Uniformity of rational points*, *J. Amer. Math. Soc.*, to appear.
- [D1] L. Denis, *Géométrie Diophantienne sur les Modules de Drinfeld*, in *The Arithmetic of function fields*, D. Goss et al., eds., de Gruyter, Berlin, 1992, pp 285-302.
- [D2] L. Denis, *Le Théorème de Fermat-Goss*, *Trans. Amer. Math. Soc.*, **343** (1994) 713-726.
- [F1] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*. *Invent. Math.* 73, 349-366 (1983).
- [F2] G. Faltings, *Some historical notes*, in *Arithmetic Geometry*, G. Cornell, J. Silverman, eds., Springer, New York, 1986, pp. 1-8.
- [F3] G. Faltings, *Diophantine approximation on abelian varieties*, *Ann. Math.* **133** (1991) 549-576.
- [F4] G. Faltings, *The general case of S. Lang's conjecture*, Barsotti Symposium in Algebraic Geometry (Abano Terme, 1991), *Perspectives in Math.* **15**, Acad. Press, San Diego, 1994, pp. 175-182.
- [Gra] H. Grauert, *Mordells Vermutung über Punkte auf algebraischen Kurven und Funktionenkörper*. *I.H.E.S. Publ. Math.* 25, 131-149 (1965).
- [Gre] M. J. Greenberg, *Lectures on forms of higher degree*, Benjamin, New York, 1969.

- [H] E. Hrushovski, *The Mordell-Lang conjecture for function fields*, J. Amer. Math. Soc., **9** (1996) 667-690.
- [K] N. M. Katz, *Algebraic solutions of differential equations (p -curvature and the Hodge filtration)*, Inventiones Math. **18**, (1972) 1-118.
- [Ki] M. Kim, *Geometric height inequalities and the Kodaira-Spencer map*, Compositio Math., to appear.
- [L1] S. Lang, *Fundamentals of Diophantine Geometry*. Springer, New York 1983.
- [L2] S. Lang, *Number Theory III: Diophantine Geometry*, Encyclopaedia Math. Sci. **60**, Springer, Berlin 1991.
- [LdM] A. Lasjaunias and B. de Mathan, *Thue's theorem in positive characteristic* Crelle **473** (1996), 195-206.
- [Ma] Yu. I. Manin, *Rational points on an algebraic curve over function fields*. Transl. Am. Math. Soc. II Ser. 50, p. 189-234 (1966). (Russian original: Izv. Acad. Nauk U.S.S.R. 1963.) and Letter to the editor. Math. USSR Izv. 34, 465-466 (1990).
- [MDLM] M. Martin-Deschamps and R. Lewin-Ménégaux, *Applications rationnelles séparables dominantes sur une variété du type général*, Bull. Soc. Math. France **106** (1978), 279-287.
- [Ms] R. Mason, *Diophantine equations over function fields*, London Math. Soc. Lecture Notes **96**, Cambridge Univ. Press 1984.
- [dM] B. de Mathan, *Approximation exponents for algebraic functions in positive characteristic*, Acta Arith. **LX** (1992), 359-370.
- [MR] W. H. Mills and D. P. Robbins, *Continued fractions of certain algebraic power series*, J. Number Theory **23** (1986), 388-404.

- [Mi] J. S. Milne, *Arithmetic duality theorems*, Academic Press, Orlando, 1986.
- [MB] L. Moret-Bailly, *Pinceaux de variétés abéliennes*, Asterisque **129** (1985).
- [Mo] A. Moriwaki, *Height inequality of non-isotrivial curves over function fields*, J. of Algebraic Geometry, **3** (1994) 249-264.
- [Og] A. Ogus, *F-crystals and Griffiths transversality*, in Proc. Int. Symp. on Algebraic Geometry, Kyoto, 1977. Kinokuniya, Tokyo, 1978.
- [O] C. F. Osgood, *Effective bounds on the "diophantine approximation" of algebraic functions over fields of arbitrary characteristics and applications to differential equations*, Indag. Math. **37**, (1975), 105- 119.
- [Ph] T. Pheidas, *Hilbert's Tenth Problem for fields of rational functions over finite fields*, Invent. Math., **103** (1991), 1-8.
- [Sa1] P. Samuel, *Lectures on old and new results on Algebraic curves*. Bombay: Tata Inst. Fund. Res. 1966.
- [Sa2] P. Samuel, *Compléments à un article de Hans Grauert sur la conjecture de Mordell*. I.H.E.S. Publ. Math. no. 29 (1966) 55-62.
- [Si] J. H. Silverman, *Arithmetic distance functions and height functions in Diophantine geometry*, Math. Ann. **279** (1987) 193-216.
- [Sz] L. Szpiro, *Seminaire sur les pinceaux de courbes de genre au moins deux*, Asterisque **86** (1981).
- [Vj] P. Vojta, *Diophantine approximations and value distribution theory*, Lecture Notes in Math. **1239**, Springer, Berlin 1987.
- [V1] J. F. Voloch, *Diophantine approximation in positive characteristic*, Periodica Math. Hungarica **19** (1988), 217-225.
- [V2] J. F. Voloch, *On the conjectures of Mordell and Lang in positive characteristic*, Invent. Math. **104** (1991) 643-646.

[V3] J. F. Voloch, *A diophantine problem on algebraic curves over function fields of positive characteristic*, Bull. Soc. Math. France **119** (1991) 121-126.

[V4] J. F. Voloch, *Explicit p -descent for elliptic curves in characteristic p* , Compositio Math. **74** (1990) 247-258.

[V5] J. F. Voloch, *Diophantine Approximation on Abelian varieties in characteristic p* , Amer. J. Math., **117** (1995), 1089-1095.

[V6] J. F. Voloch, *Diophantine approximation in characteristic p* Monatshefte für Mathematik, **119** (1995), 321-325.

[W] J. T.Y. Wang, *S -integral points of projective space omitting hyperplanes over function fields of positive characteristic*, preprint, 1995.

Dept. of Mathematics, Univ. of Texas, Austin, TX 78712, USA

e-mail: voloch@math.utexas.edu