Non-periodic long-range order for one-dimensional pair interactions

Aernout C. D. van Enter
Institute for Theoretical Physics
Rijksuniversiteit Groningen
P.O. Box 800
9747 AG Groningen
The Netherlands
AENTER@TH.RUG.NL

Boguslaw Zegarlinski
Department of Mathematics
Imperial College
London
United Kingdom
boz@ic.ac.uk

April 25, 1997

Abstract

We show how in one dimension non-periodic (quasiperiodic) long-range order can arise due to long-range translation-invariant pair interactions with quasiperiodically alternating signs. We discuss the Parisi overlap distribution between the uncountably many pure states.

1 Introduction

Since the discovery of quasicrystals [27] the description of systems which are ordered in a non-periodic manner, either quasiperiodically or even more irregularly, has been a topic of considerable interest and physical relevance, see for example [12, 11, 18, 26] and the references mentioned there. Moreover, the study of such irregular types of long-range order may be of use in understanding some conceptual aspects of the spin-glass problem.

One question of interest is to show the possibility that such a non-periodic long-range order arises for spin models with translation-invariant interactions, in which the spins can take a finite number of values. At zero temperature this has been achieved for models of spatial dimension at least two, mostly on a lattice, with nearest neighbour interactions, based on the theory of non-periodic tilings. In one-dimensional models, one needs infinite-range interactions [23], which however can decay arbitrarily fast and can be chosen to be of finite-body type [9]. At positive temperatures up to now there have been only limited results. In one dimension to obtain any type of long-range order, the interactions are necessarily of infinite range, and, moreover, they have to decay rather slowly. In [6] it was shown that for some long-range four-body interaction, of a rather artificial form, long-range order of Thue-Morse type can occur. A related argument for pair interactions with huge gaps (that is, couplings are non-zero only for pairs of spins at exceptional distances from each other) was also indicated. Here we use similar ideas to show the possibility of *a quasiperiodic long-range order occurring for a translation-invariant pair interaction of alternating sign*, in such a way that the sign of the correlation function essentially follows the sign of the interaction. In fact, our argument is in some sense inverted, in that the sign of the wished-for interaction follows the sign of some quasiperiodic correlation function.

Long-range translation-invariant pair interactions with alternating sign are physically much more realistic than the interactions considered in [6]. One of the best-known examples is the RKKY-interaction which plays an important role in the modelling of spin-glasses. For a recent review on results for long-range interactions we refer to [28]. We remark that a system with non-periodic long-range order has uncountably infinitely many pure states, a property which often is suggested to be characteristic for spin-glasses. (For a further discussion about analogies between non-periodic systems and spin-glasses, we refer to [5, 7, 28].) The main difference seems to be the absence of disorder; however, there have also been attempts to describe spin-glass behaviour with non-random models [4, 15, 16, 2, 3, 24].

As a side remark we observe that it is much simpler to obtain a non-periodic (helical) long-range order for continuous vector spin models (compare [25], p.163, note 26). One can consider, for example, a rotator model with Hamiltonian

$$H = -\Sigma_{i,j} J(i-j) cos(\theta_i - \theta_j - (i-j) \times \alpha) \tag{1.1}$$

for α an irrational multiple of 2π, which is equivalent by a simple change of variables to a ferromagnetic model if the $J(i-j)$ are positive. The ferromagnetic long-range order of the transformed model is equivalent to a non-periodic long-range order in the original model.

2 The result

In [28] it was conjectured that under reasonable conditions an Ising pair interaction whose Fourier transform has its maximum at an irrational point should show non-periodic long-range order. It can easily be seen, as is also noticed there, that arguments implying that the behaviour of the Fourier transform of the interaction function governs the nature of the phase transition cannot be valid in complete generality. Here we show some results which at least go in the right direction towards confirming this picture.

We consider the circle $[0, 2\pi)$ and the irrational rotation map (over an irrational angle α):

$$T_\alpha \theta = \theta + \alpha \tag{2.1}$$

To any θ we associate the sequence $\sigma_\alpha(\theta) \in \{-1, +1\}^Z$ defined by $\sigma_{n,\alpha}(\theta) = 1$ if $\theta + n\alpha \pmod{2\pi} \in [\frac{-\pi}{2}, \frac{+\pi}{2})$ and $\sigma_{n,\alpha}(\theta) = -1$ if $\theta + n\alpha \pmod{2\pi} \in [\frac{\pi}{2}, \frac{3\pi}{2})$. Here α is an irrational multiple of 2π.

Then the irrational rotation corresponds to the translation map on these sequences, and the rotation invariant Lebesgue measure on the circle corresponds to a translation-invariant, ergodic measure μ_α which is induced on the space $\{-1, +1\}^Z$ by the map $\theta \to \sigma_\alpha(\theta)$. It is straightforward to compute the pair correlation functions of these measures, and to see that they have a non-periodic but quasiperiodic long-range order. Indeed, let

$$\mu_\alpha(\sigma_0 \sigma_n) = f_\alpha(n) \tag{2.2}$$

with

$$f_\alpha(n) = 1 - |\frac{((n \times \alpha)\ (mod\ 2\pi))}{\pi}| \tag{2.3}$$

then the $f_\alpha(n)$ form a quasiperiodic sequence of pair correlation functions. (The Fourier transform of these pair correlations lives on a dense set of points in the interval $[0, 2\pi)$, given by the integer multiples of α (mod 2π)). It holds that $sgn(f_\alpha(n)) = sgn(cos n\alpha)$.

We will apply the Israel-Bishop-Phelps theorem [1, 13], [14] Ch.V, [10] Ch.16, to conclude the existence of a similar long-range order for an appropriate Gibbs measure on the configuration-space (space of sequences) $\{-1, +1\}^Z$. For the theory of Gibbs measures for lattice models, we refer to [8, 10, 14]. Applying the Israel-Bishop-Phelps theorem works in the same way in any dimension, for simplicity we thus consider here the one-dimensional case only. Let us consider in the following cones of translation-invariant interactions given by:

$$C_\alpha = \{J(n) : sgn(J(n)) = sgn(f_\alpha(n)),\ \Sigma_n|J(n)| \leq \infty\} \tag{2.4}$$

such that one has formal Ising Hamiltonians of the form

$$H = -\Sigma_{i,j\in Z} J(i-j)\sigma_i\sigma_j \tag{2.5}$$

The Israel-Bishop-Phelps theorem in our case reads:

Theorem 2.1 *For every irrational α and every $\epsilon \geq 0$ there exists a translation-invariant pair interaction $J_{\epsilon,\alpha} \in C_\alpha$, such that $\Sigma_n|J_{\epsilon,\alpha}(n)| \leq \frac{ln\ 2}{\epsilon}$, for which there exists a translation-invariant Gibbs measure $\mu_{J_{\epsilon,\alpha}}$ which satisfies*

$$sgn(\mu_\alpha(\sigma_0\sigma_n))[\mu_{J_{\epsilon,\alpha}}(\sigma_0\sigma_n) - \mu_\alpha(\sigma_0\sigma_n)] \geq\ -\epsilon \tag{2.6}$$

This means that there exists a Gibbs measure with the same kind of quasiperiodic long-range order in the two-point correlation functions as occurs in μ_α, while the sign of the interaction between spins at distance n coincides with the sign of the correlation function at distance n, at least when this correlation function is not too close to zero. For pair interactions this puts a limit on the unavoidable frustration. *The decomposition into extremal Gibbs measures of any such a $\mu_{J_{\epsilon,\alpha}}$ contains uncountably many elements* (for any extremal non-periodic Gibbs measure occurring in the decomposition its countably many translates occur with the same weight, which thus has to be zero; as countably many elements of measure zero still only add up to a measure zero contribution, one needs to have uncountably many extremal elements in the decomposition).

To get some more control over the cone of interactions, which gets closer to the Fourier transform having a maximum at α, at the cost of losing some control over the correlation functions, we can replace C_α by its subcone

$$C'_\alpha = \{J(n) : sgn(J(n)) = sgn(f_\alpha(n)),\ \Sigma_n \frac{J(n)}{cosn\alpha} \leq \infty\} \tag{2.7}$$

This means that in the Fourier series $\Sigma_n\ J(n)cosnx$, at the point $x = \alpha$ not only are all the terms positive, but the $J(n)$ have to be small whenever the term $cosn\alpha$ is small.

For the corresponding Gibbs measure for some element of the subcone we obtain the inequality

$$\frac{1}{cosn\alpha}[\mu_{J_{\epsilon,\alpha}}(\sigma_0\sigma_n) - \mu_\alpha(\sigma_0\sigma_n)] \geq -\epsilon \tag{2.8}$$

which still implies that non-periodic long-range order occurs.

Remark: Similarly to the case of the Fibonacci system [5], it is possible to compute the Parisi overlap distribution $p(q)$ of the μ_α. Let us recall that the overlap between two configurations $\sigma^{\mathbf{1}}$ and $\sigma^{\mathbf{2}}$ is given by

$$q = lim_{N \to \infty} \frac{1}{N} \Sigma_{n=1,\ldots N} \ \sigma_n^1 \sigma_n^2 \tag{2.9}$$

Its distribution with respect to the product-measure $\mu_\alpha \otimes \mu_\alpha$ is the Parisi overlap distribution which plays an important role in the mean-field theory of spin-glasses [17]. In our case it is given by

$$p(q) \ dq = \frac{1}{2} 1_{[-1,+1]} dq \tag{2.10}$$

which, as one sees, is non-trivial, continuous, and is the same for each α. This behaviour for the overlap distribution, as well as the earlier results of [5], illustrates the point emphasised in [19, 20, 21], that (also non-trivial) overlap-distributions in great generality are self-averaging. Here the self-averaging property is trivially fulfilled–there is no averaging to be done–but the overlap distribution can be explicitly computed and shown to be non-trivial.

3 Comments and Conclusions

We used the Israel-Bishop-Phelps theorem to show the occurrence of quasiperiodic long-range order at positive temperatures for translation-invariant pair interactions of (quasiperiodically) alternating signs. As usual the problem with applying the Israel-Bishop-Phelps theorem is its non-constructive character. One knows that there exists an interaction within a certain class of interactions whose Gibbs states give rise to some prescribed type of long-range order, but one does not know this interaction precisely. However, in cases where there is no reflection positivity available, and also no contour argument has been found, this is essentially the only rigorous method available to us. It should be helpful for a further progress in this domain to see that *the phase transition with non-periodic long-range order (with uncountably many pure Gibbs phases) can occur for a certain type of interaction.* As compared to the earlier results of [6], the class of interactions we consider contains *physically much more realistic potentials.* Moreover our results for the first time *link the Israel-Bishop-Phelps theorem with classes of interactions whose Fourier transforms have certain prescribed properties.* Attempts to read off the behaviour of the long-range correlation functions and the phase transition from the properties of the Fourier transform of an interaction are of course widespread; they form the foundation of spin-wave theory. A rigorous version of these arguments holds in fact in the mean-field limit [22], but for models with interactions of shorter range it is unclear under which conditions such arguments can be justified. Our results support the validity of some of such considerations, if only in a rather implicit fashion.

Similarly to what was observed in [5, 7, 28], we find that there is a suggestive resemblance between the properties of a non-periodically ordered system and the presumed properties of spin-glass models.

Acknowledgements: A.C.D.v.E. thanks Imperial College for its hospitality. His interest in non-periodic long-range order, as well as some of the ideas used here, owe much to earlier collaborations and discussions with Bert Hof, Jacek Miękisz and Charles Radin.

This research has been supported by EU contract CHRX-CT93-0411.

References

[1] E. Bishop and R. R. Phelps. The support functionals of a convex set. AMS Proceedings of Symposia in Pure Mathematics, Vol. 7:25-35, Amer. Math. Soc., Providence, R.I. 1963.

[2] J.-P. Bouchaud and M. Mézard. Self-induced quenched disorder, a model for the glass transition. *J. Pysique* I, 4:1109–1114, 1994.

[3] C. F. Cugliandolo, J. Kurchan, G. Parisi and F. Ritort. Matrix models as solvable glass models. *Phys. Rev. Lett.* 74:1012–1015, 1995.

[4] T. Eisele. Equilibrium and nonequilibrium theory of a geometric long-range spin glass. Les Houches summer school 1984 on: Critical Phenomena, Random Systems, Gauge Theories, Eds. K. Osterwalder and R. Stora, Vol II:1155–1165, 1986.

[5] A. C. D. van Enter, A. Hof and J. Miękisz. Overlap distributions for deterministic systems with many pure states. *J. Phys. A, Math. Gen.*, 25: L1133–1137, 1992.

[6] A. C. D. van Enter and J. Miękisz. Breaking of periodicity at positive temperatures. *Comm. Math. Phys.*, 134:647–651, 1990.

[7] A. C. D. van Enter. On the set of pure states for some systems with non-periodic long-range order. *Physica A*, 232:600–607, 1996.

[8] A. C. D. van Enter, R. Fernández and A. D. Sokal. Regularity properties and pathologies of position-space renormalization-group transformations:scope and limitations of Gibbsian theory. *J. Stat. Phys.*, 72:879–1167, 1993.

[9] C. Gardner, J. Miękisz, A. C. D. van Enter and C. Radin. Fractal symmetry in an Ising model. *J.Phys.A, Math.Gen.*, 22:L1019–1022, 1989.

[10] H.-O. Georgii. *Gibbs Measures and Phase Transitions.* Walter de Gruyter (de Gruyter Studies in Mathematics, Vol. 9), Berlin–New York, 1988.

[11] A. Hof. *Quasicrystals, Aperiodicity and Lattice Systems.* Ph. D. Thesis, Rijksuniversiteit Groningen, 1992.

[12] *Beyond Quasicrystals (les Houches 1994)*, Eds. F. Axel and D. Gratias. Springer, Berlin etc., 1995.

[13] R. B. Israel. Existence of phase transitions for long-range interactions. *Comm. Math. Phys.*, 43:59–68, 1975.

[14] R. B. Israel. *Convexity in the theory of lattice gases.* Princeton University Press, Princeton, N.J. 1979.

[15] E. Marinari, G. Parisi and F. Ritort. Replica field theory for deterministic models(I): Binary sequences with low autocorrelation. *J. Phys. A, Math. Gen.*, 27:7615–7645, 1994.

[16] E. Marinari, G. Parisi and F. Ritort. Replica field theory for deterministic models(II): A non-random spin glass with glassy behaviour. *J.Phys. A, Math. Gen.*, 27:7647–7668, 1994.

[17] M. Mézard, G. Parisi and M. A. Virasoro. *Spin Glass Theory and Beyond.* World Scientific, Singapore, 1987.

[18] J. Miękisz. *Quasicrystals. Microscopic models of non-periodic structures.* Leuven Lecture Notes in Mathematical and Theoretical Physics, Vol. 5, 1993.

[19] C. M. Newman and D. L. Stein. Non-mean-field behavior of realistic spin glasses. *Phys. Rev. Lett.*, 76:515–518.

[20] C. M. Newman. Topics in disordered systems. ETH Lecture Notes, Birkhäuser, Boston, to appear, 1997.

[21] C. M. Newman and D. L. Stein. Thermodynamic chaos and the structure of short-range spin glasses, in Mathematics of Spin glasses and neural networks, Eds. A. Bovier and P. Picco, Birkhäuser, Boston, to appear, 1997.

[22] C. Pisani and C. J. Thompson. Generalized classical theory of magnetism. *J. Stat. Phys.*, 46:971–982, 1987.

[23] C. Radin and L. S. Schulman. Periodicity and classical ground states. *Phys. Rev. Lett.*, 51:621–622, 1983.

[24] F. Ritort. Glassyness in a model without energy barriers. *Phys. Rev. Lett.* 75:1190–1193, 1995.

[25] D. Ruelle. *Statistical mechanics: rigorous results.* W. A. Benjamin, Reading, Massachusetts, 1969.

[26] M. Senechal. *Quasicrystals and geometry.* Cambridge University Press, 1995.

[27] D. Schechtman, I. Blech, D. Gratias and J. W. Cahn. Metallic phase with long-range orientational order and no translational symmetry. *Phys. Rev. Lett.*, 53:1951–1953, 1984.

[28] B. Zegarlinski. Spin systems with long-range interactions. *Rev. Math. Phys.*, 6:115–134, 1994.