

THE UNIVERSITY OF TEXAS AT AUSTIN  
DEPARTMENT OF MATHEMATICS

The Preliminary Examination in Probability  
Part I

Fri, Jan 12, 2024

**Problem 1.** Let  $T : [0, 1] \rightarrow [0, 1]$  be given by

$$T(0) = 0 \text{ and } T(x) = \{1/x\} \text{ for } x \in (0, 1],$$

where  $\{a\} := \sup\{a - m : m \in \mathbb{Z} \text{ and } m \leq a\} = a - \lfloor a \rfloor$  is the fractional part of  $a$ . Show that  $T_*\mu = \mu$ , where  $T_*\mu$  is the pushforward of  $\mu$  via  $T$  and

$$\mu(B) = \int_B \frac{1}{1+x} dx \text{ for } B \in \mathcal{B}([0, 1]).$$

(Note:  $T$  is (one of several things) known as the **Gauss map**.)

**Problem 2.** Let  $\mu$  be a probability measure on  $\mathbb{R}$ , and let  $\varphi$  be its characteristic function. Show that  $\mu$  is diffuse (has no atoms) if

$$\lim_{t \rightarrow \infty} |\varphi(t)| = \lim_{t \rightarrow -\infty} |\varphi(t)| = 0.$$

(Hint: For  $a \in \mathbb{R}$ , compute  $\lim_{T \rightarrow \infty} \int_{-T}^T e^{-ita} \varphi(t) dt$ .)

**Problem 3.** Given  $X \in \mathbb{L}^2(\mathcal{F})$  and two sub- $\sigma$ -algebras  $\mathcal{G}, \mathcal{H}$  of  $\mathcal{F}$  such that  $\mathcal{G} \subseteq \mathcal{H}$ , show that

$$\mathbb{E}[\text{Var}[X | \mathcal{G}]] \geq \mathbb{E}[\text{Var}[X | \mathcal{H}]],$$

where  $\text{Var}[X | \mathcal{K}] := \mathbb{E}[(X - \mathbb{E}[X | \mathcal{K}])^2 | \mathcal{K}]$  for  $\mathcal{K} \subseteq \mathcal{F}$ . When does the equality hold?

(Note:  $\text{Var}[X | \mathcal{K}]$  is called the **conditional variance** of  $X$  given  $\mathcal{K}$ .)